


**ANNUAL REPORT
ON
GYNECOLOGIC ONCOLOGY
2007**



**DIVISION OF GYNECOLOGIC ONCOLOGY
DEPARTMENT OF OBSTETRICS AND GYNECOLOGY
FACULTY OF MEDICINE, CHIANG MAI UNIVERSITY
CHIANG MAI, THAILAND**

ANNUAL REPORT 2007 GYNECOLOGIC ONCOLOGY

**DIVISION OF GYNECOLOGIC ONCOLOGY
DEPARTMENT OF OBSTETRICS AND GYNECOLOGY
FACULTY OF MEDICINE, CHIANG MAI UNIVERSITY
CHIANG MAI, THAILAND**

WEBSITE : <http://www.med.cmu.ac.th/dept/obgyn/Unit/onco/oncofront.htm>

GYNECOLOGIC ONCOLOGY STAFFS 2007

Professor Jatupol Srisomboon, M.D.
Associate Professor Prapaporn Suprasert, M.D.
Assistant Professor Chailert Phongnarisorn, M.D.
Assistant Professor Chalong Cheewakriangkrai, M.D.
Assistant Professor Kittipat Charoenkwan, M.D.
Assistant Professor Sitthicha Siriaree, M.D.
Charuwan Tantipalakorn, M.D.
Chumnan Kietpeerakool, M.D.
Sukunya Yanunto, MSc.
Narisa Sribanditmongkol, MSc.
Aree Pantusart, BSc.
Tosapol Chainoy, B.A.

รายงานประจำปี 2550

หน่วยมะเร็งวิทยานรีเวช
ภาควิชาสูติศาสตร์และนรีเวชวิทยา
คณะแพทยศาสตร์ มหาวิทยาลัยเชียงใหม่

อนุสาขามะเร็งวิทยานรีเวช

ศาสตราจารย์ นายแพทย์ จตุพล ศรีสมบูรณ์
รองศาสตราจารย์ แพทย์หญิง ประภาพร สู่ประเสริฐ
ผู้ช่วยศาสตราจารย์ นายแพทย์ ชัยเลิศ พงษ์นริศ
ผู้ช่วยศาสตราจารย์ นายแพทย์ ฉลอง ชิวเกียรียงไกร
ผู้ช่วยศาสตราจารย์ นายแพทย์ กิตติภักดิ์ เจริญขวัญ
ผู้ช่วยศาสตราจารย์ นายแพทย์ สิทธิธา สิริอารีย์
อาจารย์ แพทย์หญิง จารุวรรณ ตันติพลากร
อาจารย์ นายแพทย์ ชำนาญ เกียรติพิรกุล
คุณสุกัญญา ยานันโต
คุณนริสา ศรีบัณฑิตมงคล
คุณอารีย์ พันธุศาสตร์
คุณทศพล ไชยน้อย

PREFACE

This Annual Report on Gynecologic Oncology 2007 was carried out with great contribution from Assoc. Prof. Prapaporn Suprasert, Chief of the Gynecologic Oncology Division, Khun Aree Pantusart, our ex-research nurse who resigned to take care of her daughter in Bangkok, Khun Sukunya, Khun Narisa, our new research nurses, Khun Tosapol, our research assistant, and our colleagues in the OB & GYN Department. This report also provided the accumulated data of our 11-year work in the field of gynecologic oncology.

In my opinion, the Division has made satisfactory progression and improvement in the service, academic activity, and research in this field. Numerous original articles were accepted and published in the international journals. Many staffs were invited as guest speakers in the Annual OB & GYN meetings, Thai Gynecologic Cancer Society meeting and the International Symposium on Radical Hysterectomy Dedicated to Hidekazu Okabayashi in Kyoto, Japan. A “Comprehensive Course on Cervical Cancer Prevention and Treatment” was organized by the Division with great success as more than 80 participants nationwide attending the meeting.

I sincerely appreciate Dr. Prapaporn and her colleagues for their dedication and contribution to the Gynecologic Oncology Division.

Jatupol Srisomboon, M.D
Professor & Chairman
Department of Obstetrics & Gynecology
Faculty of Medicine, Chiang Mai University
Chiang Mai 50200, Thailand
E mail: jsrisomb@mail.med.cmu.ac.th

PREFACE

This Annual Report 2007 is the eleventh volume of our work in gynecologic oncology. We served about 750 gynecologic cancer patients in 2007 which was slightly less than those in the previous years. Cervical cancer is still the leading cancer followed by ovarian and uterine cancers.

About 130 Wertheim operations were performed in our hospital. Of these, 11 cases were carried out via laparoscopic approach. With the routine nerve - sparing procedure, the morbidity especially dysfunction of bladder from Wertheim operations were decreased. In 2007, we treated some rare diseases such as primitive neuroectodermal tumor of corpus, malignant spindle tumor of vulva, 3 primary gynecologic cancer, uterine rupture from invasive mole and all of them had favorable outcome. Thirteen original studies were published in the peer-reviewed journals in 2007. Our staffs were invited as great speakers in both local and international gynecologic oncology conferences.

This report is divided into 2 sections. The first section provides the statistics of all gynecologic cancer patients in the year 2007 with accumulated data since 1997. Section II presents the infrastructure, diagnostic procedures and operations in gynecologic cancer, abstracts of the publications and presentations of our work and related specialties.

In 2007, I would like to welcome 2 new research nurses, Khun Sukunya Yanunto and Khun Narisa Sribanditmongkol and 1 new research assistant, Khun Tosapol Chainoy. They will work with us instead of Khun Aree who resigned from our Department to take care her daughter. However, I would like to thank her for preparing some part of this annual report. I take this opportunity to appreciate my colleagues and fellows for their perseverance and dedication. Finally, a special word of thankfulness goes to our Head Department of OB&GYN, Professor Jatupol Srisomboon for his incessant support.

Prapaporn Suprasert, M.D.
Associate Professor and Chief
Division of Gynecologic Oncology
Department of Obstetrics & Gynecology
Faculty of Medicine, Chiang Mai University
Chiang Mai 50200, Thailand
E-mail: psuprase@mail.med.cmu.ac.th

CONTENT

	Page
SECTION I :	
➤ Gynecologic Oncology Registry, Chiang Mai 2007	3
➤ Operations and Procedures in Gynecologic Oncology	5
➤ Organ Specific Gynecologic Cancer	
Cancer of The Cervix	7
Cancer of The Ovary	13
Cancer of The Uterine Corpus	19
Cancer of The Vulva	23
Cancer of The Vagina	28
Cancer of The Fallopian Tube	31
Cancer of The Peritoneum	35
Cancer of Multiple Primaries Gynecologic Organ	41
Gestational Trophoblastic Disease	45
SECTION II :	
➤ Medical Personnel and Facilities	49
➤ Diagnostic Procedures	52
& Gynecologic Oncology Operations	
➤ Publications & Presentations	53

SECTION I

- **Gynecologic Oncology Registry**
Chiang Mai University : 2007

- **Operations and Procedures**
In Gynecologic Oncology

- **Organ Specific Gynecologic Cancer**
 - Cancer of The Cervix
 - Cancer of The Ovary
 - Cancer of The Uterine Corpus
 - Cancer of The Vulva
 - Cancer of The Vagina
 - Cancer of The Fallopian Tube
 - Cancer of The Peritoneum
 - Cancer of Multiple Primary Gynecologic Organ
 - Gestational Trophoblastic Disease

TABLE 1 : Gynecologic Oncology Registry :Chiang Mai University 1997-2007

Site	1997 Number(%)	1998 Number(%)	1999 Number(%)	2000 Number(%)	2001 Number(%)	2002 Number(%)	2003 Number(%)	2004 Number(%)	2005 Number(%)	2006 Number(%)	2007 Number(%)
Cervix	547 (75.3)	483 (72.9)	497 (75.3)	502 (71.3)	500 (70.8)	521 (69.7)	624 (71.7)	532 (66.9)	525 (66.4)	488 (66.8)	480(63.6)
Ovary	87 (12.0)	83 (12.5)	82 (12.4)	96 (13.6)	90 (12.7)	110 (14.7)	111 (12.8)	126 (15.8)	121 (15.3)	114 (15.6)	132(17.5)
Corpus	48 (6.6)	47 (7.1)	49 (7.4)	56 (8.0)	63 (8.9)	61 (8.2)	67 (7.7)	89 (11.2)	97 (12.3)	84 (11.5)	91(12.0)
Vulva	20 (2.7)	21 (3.2)	15 (2.2)	29 (4.1)	23 (3.3)	25 (3.3)	29 (3.3)	22 (2.8)	19 (2.4)	15 (2.1)	11(1.5)
Vagina	11 (1.4)	10 (1.5)	3 (0.5)	2 (0.3)	9 (1.3)	6 (0.8)	12 (1.4)	5 (0.6)	4 (0.5)	5 (0.7)	6(0.7)
FT	-	2 (0.3)	3 (0.5)	5 (0.7)	3 (0.4)	4 (0.5)	6 (0.7)	5 (0.6)	4 (0.5)	7 (1.0)	7(0.9)
PPA	-	-	2 (0.3)	1 (0.1)	-	2 (0.3)	7 (0.8)	3 (0.4)	4 (0.5)	6 (0.8)	11(1.5)
GTT	14 (1.9)	16 (2.4)	8 (1.2)	13 (1.9)	18 (2.6)	19 (2.5)	14 (1.6)	13 (1.6)	17 (2.1)	12 (1.6)	17(2.3)
Total	727(100)	662(100)	660(100)	704(100)	706(100)	748(100)	870(100)	795(100)	791 (100)	731 (100)	755(100)

PPA = Primary Peritoneal Adenocarcinoma FT = Fallopian tube GTT = Gestational Trophoblastic Tumors

Operations and Procedures in Gynecologic Oncology

Operations and Procedures	1997 Number	1998 Number	1999 Number	2000 Number	2001 Number	2002 Number	2003 Number	2004 Number	2005 Number	2006 Number	2007 Number
Surgery for Ovarian&FT&PPA CA.	64	43	64	70	45	69	88	79	80	111	89
Surgery for Corpus CA.	33	28	26	36	43	39	47	60	75	53	80
Surgery for Vulvar CA.	10	14	5	19	12	14	21	19	14	12	8
Total Laparoscopic Hysterectomy	-	-	-	-	-	-	10	11	9	4	4
Radical Hysterectomy	55	77	113	120	116	135	150	151	149	143	120
Laparoscopic Radical Hysterectomy	-	-	-	-	-	-	-	4	18	21	11
Radical Parametrectomy	2	2	1	1	1	3	4	1	1	2	1
Laparoscopic Radical Parametrectomy	-	-	-	-	-	-	-	1	1	3	-
Simple Hysterectomy	118	110	155	182	121	89	43	35	52	55	47
Conization	66	65	79	13	14	22	16	9	10	5	15
LEEP	61	35	166	207	194	221	380	276	261	309	317
Cryosurgery	20	15	18	8	4	3	1	-	2	-	-
Colposcopy	227	235	463	371	369	306	357	399	499	627	519

Cancer of The Cervix

➤ **Distribution by**

- Age
- Gravida
- Stage and Substage
- HIV Status
- Histological Type
- Treatment

TABLE 2 : Cancer of The Cervix : Age Distribution.

Age	Number	Percent
20 – 30	6	1.3
31 – 40	74	15.4
41 – 50	172	35.8
51 – 60	135	28.1
61 – 70	47	9.8
71 – 80	40	8.3
81 – 90	5	1.0
91 – 100	1	0.2
Total	480	100

Minimum age = 28 year, Maximum age = 92 year,

Mean age = **51** year

TABLE 3 : Cancer of The Cervix : Gravidity Distribution.

Gravida	Number	Percent
0	19	3.9
1	83	17.4
2	168	35.3
3	78	16.4
4	53	11.1
>4	75	15.9
Total	476*	100

Minimum gravidity = 0, Maximum gravidity = 15,

Mean = 2.9

*Data not available = 4 cases

TABLE 4 : Cancer of The Cervix: Stage Distribution.

Stage	Number	Percent
I	185	38.5
II	130	27.1
III	138	28.8
IV	27	5.6
Total	480	100

TABLE 5 : Cancer of The Cervix: Stage and Substage Distribution.

	Stage	Number	Percent
I	Ia1	46	9.6
	Ia2	13	2.7
	Ib1	97	20.2
	Ib2	29	6.0
II	II a	26	5.4
	II b	104	21.7
III	III a	1	0.2
	III b	137	28.5
IV	IV a	10	2.1
	IV b	17	3.5
Total		480	100

Table 6 : HIV Status in Cervical Cancer Patients.

Stage	Number Negative (%)	Number Positive HIV(%)	Unknown*(%)	Total
Ia1	40 (8.3)	2 (0.4)	4 (0.8)	46 (9.6)
Ia2	10 (2.1)	3 (0.6)	-	13 (2.7)
Ib1	93 (19.4)	2 (0.4)	2 (0.4)	97 (20.2)
Ib2	29 (6.0)	-	-	29(6.0)
IIa	24 (5.0)	1 (0.2)	1 (0.2)	26 (5.4)
IIb	87 (18.1)	7 (1.5)	10 (2.1)	104 (21.7)
IIIa	-	-	1 (0.2)	1 (0.2)
IIIb	104(21.7)	10 (2.1)	23 (4.8)	137 (28.5)
IVa	8 (1.7)	-	2 (0.4)	10(2.1)
IVb	15(3.1)	-	2 (0.4)	17 (3.5)
Total	410 (85.4)	25 (5.2)	45 (9.4)	480(100)

* Unknown = Data not available

Table 7 : Cancer of The Cervix : Distribution by Histological Type.

Histological Type	Number	Percent
Squamous cell CA.	396	82.8
Well differentiated	45	9.4
Moderately differentiated	193	40.2
Poorly differentiated	66	14.1
Not define differentiated	92	19.1
AdenoCA.	69	14.1
AdenosquamousCA.	1	0.2
Neuroendocrine tumor	3	0.6
Clear cell CA.	2	0.4
Malignant lymphoma	1	0.2
HGSIL,CIS	2	0.4
Unknown	6	1.3
Total	480	100

Table 8 : Treatment of Cancer of The Cervix.

Treatment	Number	Percent
Surgery alone	110	22.9
RH & BPL	68	14.2
LRHPL	8	1.7
Extended hysterectomy	17	3.5
TAH	15	3.1
Laparoscopic radical trachelectomy & BPL	1	0.2
RP & BPL	1	0.2
Chemotherapy	16	3.3
Radiation ± Chemotherapy	282	58.8
Combined treatment	71	14.8
TAH + CT	1	0.2
TAH + RT	1	0.2
TAH + Brachytherapy	1	0.2
RH + Brachytherapy	2	0.4
RH + RT	3	0.6
RH + CCRT	37	7.7
RH + CCRT + Brachytherapy	1	0.2
RH + CT	7	1.4
Extended hysterectomy + CCRT	1	0.2
NAC + RH	2	0.4
NAC + RH + CCRT	3	0.6
LRHPL+ CCRT	1	0.2
LRHPL + Brachytherapy	2	0.4
Pelvic node dissection (abandoned RH) + CCRT	8	1.7
Symptomatic & Supportive Treatment	1	0.2
Total	480	100

RH = Radical Hysterectomy

RP = Radical Parametrectomy

TAH = Total Abdominal Hysterectomy (Extrafascial Hysterectomy)

BPL = Bilateral Pelvic Lymphadenectomy

LRHPL = Laparoscopic Radical Hysterectomy and pelvic lymphadenectomy

NAC = Neoadjuvant chemotherapy

RT = Radiation Therapy

CT = Chemotherapy

CCRT = Concurrent Chemoradiation

N.B. Number of Radical Hysterectomy & BPL = 134 cases

Cancer of The Ovary

➤ Distribution by

- Age
- Gravidity
- Marital Status
- Histology
- Histology Subtype
 - Epithelial Group
 - Germ Cell Tumor Group
 - Sex cord-stromal Group
 - Others Group
- Stage
 - Epithelial Group
 - Germ Cell Group
 - Sex cord-stromal Group
 - Other Group
- Stage and Histology
- Treatment

TABLE 9 : Cancer of The Ovary : Age Distribution.

Age	Number	Percent
1 –10	2	1.5
11– 20	9	6.8
21 – 30	4	3.0
31 – 40	15	11.4
41 – 50	47	35.6
51 – 60	39	29.5
61 – 70	8	6.0
71 – 80	8	6.0
Total	132	100

Minimum age = 7 years., Maximum age = 77 years.,
Mean age = 46 years.

TABLE 10 : Cancer of The Ovary : Gravidity Distribution.

Gravidity	Number	Percent
Nulligravida	43	32.6
Multigravida	89	67.4
Total	132	100

TABLE 11 : Cancer of The Ovary : Marital Status Distribution.

Marital status	Number	Percent
Single	28	21.1
Married	92	69.7
Divorced	6	4.5
Widow	6	4.5
Total	132	100

TABLE 12 : Cancer of The Ovary : Histological Distribution.

Histology	Number	Percent
Coelomic Epithelium	97	73.5
Germ Cell	14	10.6
Sex cord-stromal	9	6.8
Others	6	4.5
Unknown	6	4.5
Total	132	100

TABLE 13 : Epithelial Ovarian Cancer : Histological Subtype Distribution.

Subtype	Number	Percent
Serous LMP	6	6.2
Serous adeno CA	23	17.4
Mucinous LMP	13	9.8
Mucinous adeno CA	9	6.8
Endometrioid LMP	3	2.3
Endometrioid CA	12	12.4
Clear cell CA	18	18.6
Mixed Epithelial CA	7	7.2
Malignant Brenner Cell	1	1.0
Undifferentiated CA	1	1.0
AdenoCA	4	4.1
Total	97	100

TABLE 14 : Ovarian Germ Cell Tumor (GCT) : Histological Subtype Distribution.

Subtype	Number	Percent
Dysgerm	1	7.1
Endodermal sinus tumor	4	28.6
Immature teratoma	5	35.7
Squamous cell CA arising in mature teratoma	3	21.4
Papillary thyroid CA arising in mature teratoma	1	7.1
Total	14	100

TABLE 15 : Sex cord-stromal tumor : Histological Subtype Distribution.

Subtype	Number	Percent
Adult granulosa cell tumor	1	50
Juvenile granulosa cell tumor	1	50
Total	2	100

TABLE 16 : Others : Histological Subtype Distribution.

Subtype	Number	Percent
carcinosarcoma	2	33.3
Sertoli-leydig cell tumor	1	16.7
Non-small cell neuroendocrine CA	1	16.7
Metastasis mucinous CA	2	33.3
Total	6	100

TABLE 17 : Epithelial Ovarian Cancer : Stage Distribution.

Stage	Number	Percent
I	46	47.4
II	11	11.3
III	21	21.6
IV	7	7.2
Recurrence	12	12.4
Total	97	100

TABLE 18 : Germ Cell Ovarian Cancer: Stage Distribution.

Stage	Number	Percent
I	9	64.3
II	-	-
III	2	14.3
IV	1	7.1
Recurrence	2	14.3
Total	14	100

TABLE 19 : Sex cord-stromal : Stage Distribution.

Stage	Number	Percent
I	6	66.7
II	-	-
III	1	11.1
IV	2	22.2
Total	9	100

TABLE 20 : Others : Stage Distribution.

Stage	Number	Percent
I	2	33.3
II	-	-
III	2	33.3
IV	-	-
Recurrence	1	16.7
Unknown	1	16.7
Total	6	100

TABLE 21 : Ovarian Cancer : Stage and Histology Distribution.

Stage	Epithelial	(%)	Germ Cell	(%)	Sex cord-stromal	(%)	Others	(%)
I	46	47.4	9	64.3	6	66.7	2	33.3
II	11	11.3	-	-	-	-	-	-
III	21	21.6	2	14.3	1	11.1	2	33.3
IV	7	7.2	1	7.1	-	22.2	-	-
Recurrence	12	12.4	2	14.3	2	-	1	16.7
Unknown	-	-	-	-	-	-	1	16.7
Total	97	100	14	100	9	100	6	100

TABLE 22 : Cancer of The Ovary : Primary Treatment and Adjuvant Chemotherapy.

Treatment	Without Adj.Chemo (%)	With Adj.Chemo. (%)	Total (%)
Complete SSP.	15 (11.4)	18 (13.6)	33(25)
Incomplete SSP.	8	5	13
Optimal debulking	1	4	5
Suboptimal debulking	1	6	7
Interval debulking (optimal)	-	1	1
Interval debulking (sub-optimal)	-	1	1
Biopsy	-	2	2
No surgery	6	1	7
Referred case			
With complete SSP	3	17	20
With incomplete SSP	10	26	36
With optimal debulking	-	1	1
With suboptimal debulking	1	4	5
With biopsy	-	1	1
Total	45	87	132

SSP = Surgical Staging Procedure

Neoadjuvant Chemotherapy alone (no surgery) = 7 cases

Neoadjuvant Chemotherapy followed by Optimal Debulking = 2 cases

Neoadjuvant Chemotherapy followed by Suboptimal Debulking = 2 cases

TABLE 23 : Ovarian Cancer : Outcome of Treatment.

Outcome \ Stage	I	II	III	IV	Recurrence	Unknown	Total
Complete Chemotherapy							
- Under follow-up without disease	16	7	3	1	2	-	29
- Under follow-up with partial response	-	1	4	2	1	-	8
- Under follow-up with progress or persist of disease \pm 2 nd or 3 rd line chemotherapy or recurrence	1	-	3	-	1	1	6
Incomplete Chemotherapy							
- During treatment	9	1	9	2	2	1	24
- During treatment with progress of disease	1	-	3	2	4	-	10
- Lost to follow-up with disease	1	1	1	1	1	1	6
- Hormonal treatment	-	-	-	2	1	-	3
- Dead	-	-	2	-	1	-	3
- Follow-up with S/S	1	-	1	-	2	-	4
No Chemotherapy							
- Under follow-up without disease	32	1	2	-	1	-	36
- Under follow up with disease	-	-	-	-	1	1	2
- Lost to follow up	1	-	-	-	-	-	1
Total	62	11	28	10	17	4	132

S/S = Symptomatic & Supportive Treatment

Cancer of The Uterine Corpus

➤ **Distribution by**

- Age
- Menopausal Status
- Underlying Medical Diseases
- Parity and Marital Status
- Stage
- Histology
- Treatment

Table 24 : Cancer of The Corpus : Age Distribution.

Age	Number	Percent
20 – 30	1	1.0
31 – 40	4	4.4
41 – 50	24	26.4
51 – 60	37	40.7
61 – 70	15	16.5
71 – 80	10	11.0
Total	91	100

Minimum age = 30 years., Maximum age = 78 years.,

Mean age = 56 years

Table 25 : Cancer of The Corpus: Distribution by Menopausal Status.

Menopausal Status	Number	Percent
Premenopause	28	30.8
Postmenopause*	63	69.2
Total	91	100

* 4 cases = surgical menopause

Table 26 : Cancer of The Uterine Corpus: Distribution by Underlying Medical Diseases.

Medical Disease	Number	Percent
No	52	57.1
Diabetes Mellitus (DM)	4	4.4
Hypertension (HT)	17	18.7
HT+ DM	6	6.6
HT+ Heart Disease	3	3.3
DM+HT+ Heart Disease	1	1.1
Thyrotoxicosis	1	1.1
Heart Disease	2	2.2
Other	5	5.5
Total	91	100

Table 27 : Cancer of The Uterine Corpus : Distribution by Parity.

Parity	Number	Percent
0	19	20.9
1	12	13.2
2	20	22.0
3	20	22.0
4	13	14.3
>4	7	7.7
Total	91	100

Minimum = 0, Maximum = 7

Table 28 : Cancer of The Uterine Corpus : Distribution by Marital Status.

Marital Status	Number	Percent
Single	9	9.9
Married	58	63.7
Divorced	9	9.9
Widowed	15	16.5
Total	91	100

Table 29 : Cancer of The Uterine Corpus : Distribution by Surgical Staging.

Surgical Stage	Number	Percent
I		
Ia	12	13.2
Ib	18	19.8
Ic	15	16.5
II		
IIa	3	3.3
IIb	7	7.7
III		
IIIa	5	5.5
IIIb	1	1.1
IIIc	12	13.2
IV		
IVb	8	8.8
Recurrence (referred)	7	7.7
Unknown	3	3.3
Total	91	100

Table 30 : Cancer of The Corpus : Histologic Distribution.

Histology	Number	Percent
Endometrioid Adenocarcinoma		
Grade I	36	39.6
Grade II	12	13.2
Grade III	16	17.6
Carcinosarcoma	8	8.8
Leiomyosarcoma	5	5.5
Endometrial Stromal Sarcoma	1	1.1
Clear Cell Carcinoma	2	2.2
Primitive neuroectodermal tumor (PNET)	1	1.1
Serous Adenocarcinoma	2	2.2
Undifferentiated carcinoma	1	1.1
Mixed type	6	6.6
Unknown	1	1.1
Total	91	100

Table 31 : Treatment of Corpus Cancer.

Treatment	Number	Percent
Surgery		
Complete SSP	25	27.5
Complete SSP + RT	22	24.2
Complete SSP + CT	13	14.3
Complete SSP + CT + RT	2	2.2
Complete SSP + brachytherapy	6	6.6
Incomplete SSP	6	6.6
Incomplete SSP + RT	6	6.6
Incomplete SSP + brachytherapy	1	1.1
Incomplete SSP + CT	2	2.2
Incomplete SSP + hormone	1	1.1
Incomplete SSP + CT + S&S	1	1.1
Previous vaginal hysterectomy + CT	1	1.1
Previous TAH + CT	1	1.1
Previous complete SSP +debulking tumor	1	1.1
No Surgery		
RT	2	2.2
No treatment		
	1	1.1
Total	91	100

SSP = Surgical Staging Procedure

RT = Radiation Therapy

CT = Chemotherapy

CCRT = Concurrent Chemoradiation

S&S = Supportive symptomatic treatment

TAH = Trans - abdominal hysterectomy

Table 32: Outcome of Treatment of Corpus Cancer.

Outcome	Number	Percent
- During Rx	18	19.8
- Under Follow-up without Disease	56	61.5
- Under Follow-up with Disease	4	4.4
- Lost to Follow-up with Disease	5	5.5
- Symptomatic & Supportive	3	3.3
- Dead	3	3.3
- Awaiting for treatment	1	1.1
- Data not available	1	1.1
Total	91	100

Cancer of The Vulva

➤ **Distribution by**

- Age
- Stage
- Histology
- Treatment

Table 33 : Cancer of The Vulva : Age Distribution.

Age	Number	Percent
31-40	2	18.2
41-50	1	9.1
51-60	4	36.4
61-70	2	18.2
71-80	1	9.1
80-90	1	9.1
Total	11	100

Minimum age = 31 years., Maximum age = 82 years.,

Mean age = 57 years.

Table 34 : Cancer of The Vulva : Stage Distribution.

Stage	Number	Percent
I	2	18.2
II	4	36.4
III	5	45.4
IV	-	-
Total	11	100

Table 35 : Cancer of The Vulva : Histological Type Distribution.

Histological type	Number	Percent
Squamous cell Carcinoma		
Well-differentiated	7	63.6
Moderately-differentiated	1	9.1
Squamous cell Carcinoma	1	9.1
(well differentiated) + Paget's disease		
Melanoma	1	9.1
Malignant spindle tumor	1	9.1
Total	11	100

Table 36 : Treatment of cancer of the vulva.

Treatment	Number	Percent
Surgery only	2	18.2
Radical local excision + BGND	1	9.1
Skinning vulvectomy	1	9.1
RT+CT	2	18.2
Combined treatment	6	54.5
Radical hemivulvectomy + BGND + RT	1	9.1
Radical vulvectomy + BGND + RT	1	9.1
Radical vulvectomy + BGND + TAH & BSO* +RT	1	9.1
RLE + BGND + pelvic node dissection RT	1	9.1
BGND + RT	1	9.1
WLE + RT	1	9.1
No Surgery	1	9.1
Total	11	100

- RLE = Radical local excision
WLE = Wide local excision
BGND = Bilateral groin node dissection
RT = Radiation Therapy
CT = Chemotherapy
TAH = Trans-abdominal hysterectomy
BSO = Bilateral salpingo-oophorectomy
* = Hysterectomy due to myoma

Cancer of The Vagina

➤ **Distribution by**

- Age
- Stage
- Histology
- Treatment

Table 37 : Cancer of The Vagina : Age Distribution.

Minimum age	=	1	years.
Maximum age	=	77	years.
Median age	=	59	years.
Mean age	=	53	years.

Table 38 : Cancer of The Vagina : Stage Distribution.

Stage	Number	Percent
I	1	16.7
II	1	16.7
III	3	50.0
IV	1	16.7
Total	6	100

Table 39 : Cancer of The Vagina : Histological Type Distribution.

Histological type	Number
Squamous cell CA.	
Well-differentiated.	1
Moderately-differentiated.	1
Poorly-differentiated.	1
Adenocarcinoma	
Moderately-differentiated.	1
Poorly-differentiated.	1
Endodermal sinus tumor	1
Total	6

Table 40 : Cancer of The Vagina 2006.

No	HN	Age	Stage	Histology	Treatment
1	2617128	52	III	Squamous cell CA, Poorly-differentiated.	Radiotherapy.
2	2955824	77	III	AdenoCA, Poorly- differentiated.	Lost before treatment.
3	1307750	74	III	Squamous cell CA, Well-differentiated.	Radiotherapy. (previous treatment cervical cancer stage IIB with radiotherapy last 23 years ago)
4	3014048	1	I?	Endodermal sinus tumor	Chemotherapy : BEP regimen (at pediatric oncology unit)
5	3024392	50	IV	AdenoCA, Moderate- differentiated.	Chemotherapy : PT regimen.(ovarian ,paraortic and Supraclavicular lymphnodes metastasis)
6	2993153	67	II	Squamous cell CA, Moderate-differentiated.	Radiotherapy.

BEP= bleomycin+ etoposide + cisplatin, CA = Carcinoma

Cancer of The Fallopian Tube

Table 41 : Cancer of The Fallopian Tube 2007

Data	Case 1	Case 2	Case 3
HN	2970561	2888941	2524786
Age (yrs)	52	45	70
Marital status	Married	Married	Married
Parity	3-0-0-3	2-0-0-2	3-0-0-3
Presenting symptoms	Abdominal distension	Abdominal pain , Abdominal distension	Abdominal pain
Stage	IIIC(recurrence:platinum resistant)	IC	IIA
Histology	<u>Left+Right fallopian tube</u> ; Mixed endometrioid & serous adenoCA, grade2 <u>Metastasis</u> to both ovaries, uterine serosa, omentum	<u>Right fallopian tube</u> ; endometrioid adenoCA, grade 3 <u>Metastasis</u> to external surface of appendix <u>Cytology</u> ; negative for CA	<u>Left + Right fallopian tube</u> ; serous adeno CA gr 3 <u>Metastasis</u> to corpus (myometrium)
Treatment	1.TAH+BSO (Private hospital: Aug 10,2006) 2.Chemotherapy PTx6 3.Explor lap : carcinomatosis peritonei(Province hospital: Feb 15,2006) 4.Patupilone x 5 (CMU) -> Oral etoposide x 1 -> Tamoxifen x 2 -> Weekly gemzitabine x 3 -> progression 8. Palliative treatment	1.Complete surgical staging (Dec 12,2005) 1.Chemotherapy - PT x 6 courses	1. TAH + BSO + partial omentectomy (April 4,2007) 2 Chemotherapy Carboplatin x 6 courses
Outcome	Alive with disease	Under follow-up, Disease free after 6 courses of Chemotherapy	Under follow-up, Disease free after 6 courses of Chemotherapy

Cancer of The Fallopian Tube 2007 (continue)

Case 4	Case 5	Case 6	Case 7
2989003	2994778	3026825	2831556
51	49	53	51
Married	Married	Married	Married
2-0-0-2	1-0-0-1	0	3-0-0-3
Abdominal distension	Pelvic pain	Abdominal mass	Abdominal mass
IV(lung metastasis)	IIC	IIIC	IIIC
<u>Left fallopian tube</u> ; mixed adeno CA grade 3 <u>Metastasis</u> to omentum, peritoneum <u>Cytology</u> : positive	<u>Right fallopian tube</u> ; serouscystadeno CA gr 3 confine to mucosa. <u>Metastasis</u> to right ovary	<u>Left fallopian tubes</u> ; serous adeno CA, gr 3 metastasis to paraaortic nodes <u>Right fallopian tube</u> ; carcinosarcoma	<u>Right fallopian tube</u> ; serous cystadenoCA, gr 3 <u>Metastasis</u> : to pelvic & paraaortic nodes
<u>1. Referred case</u> Omentectomy + lysis adhesion+ Lt cystectomy(April 16,2007) <u>2. Chemotherapy</u> Carboplatin x 4 -> PT x 6 → progression	<u>1.Referred case</u> TAH +BSO+omentectomy +appendectomy (April 5,2006) <u>2. Chemotherapy</u> PT x 6 -> weekly taxotere	<u>1. Complete surgical staging</u> (Oct 11,2006) <u>3. Chemotherapy</u> -PT	<u>1.Complete surgical staging</u> (Nov14,2006) <u>2. Chemotherapy</u> - PT
Palliative treatment	During treatment	During treatment	During treatment

PT = carboplatin+ paclitaxel

Cancer of The Peritoneum

Table 42 : Cancer of The Peritoneum 2007.

Data	Case 1	Case 2	Case 3
Hospital number	2959136	2954799	1641613
Age (yrs)	72	77	56
Marital status	Married	Married	Married
Parity	1-0-0-1	Unknown	2-0-0-2
Presenting symptoms	Abdominal distend, Abdominal pain	Abdominal distension	Pelvic discomfort
Stage	IIIC	IIIC	IIIC
Histology	<u>Omentum</u> : serous cystadenocarcinoma gr 3 <u>Mesenteray of large bowel</u> : serous cystadenocarcinoma gr 3	Lap. biopsy : Mucin producing adenocarcinoma	-
Treatment	1. Referred case: Explor lap to omental & mesentery of large bowel biopsy 2. Chemotherapy PT x 6 -> carboplatin x 3 -> oral etoposide	1. Carboplatin x 2 → progression 2. Supportive & symptomatic treatment	1. Chemotherapy PT x 6
Outcome	During treatment	Lost to follow up	Alive with disease, during follow- up , no symptom

Cancer of The Peritoneum 2007. (continued)

Data	Case 4	Case 5	Case 6
Hospital number	2913975	2988955	2998989
Age (yrs)	68	43	68
Marital status	Married	Married	Divorce
Parity	1-0-2-1	2-0-0-2	2-0-0-2
Presenting symptoms	Abdominal distension & chronic dry cough	Abdominal distension	Abdominal pain, Abdominal distend
Stage	IV(lung metastasis)	IIIC (recurrence)	IIIC
Histology	<u>Omentum:</u> serous cystadenocarcinoma gr 2 <u>Mesenteray & peritoneum:</u> serous cystadenocarcinoma gr 2	AdenoCA	<u>Tissue from pelvis, ileal surface:</u> carcinosarcoma <u>Both ovaries:</u> no pathology involvement
Treatment	1. Explor lap to omental & mesentery & peritoneum biopsy 28 March,2007 2. Chemotherapy: carboplatin x 6	1.TAH&BSO&partial omentectomy & peritoneal biopsy Jun,2006 (Bangkok) 2. Chemotherapy PT (carboplatin+ paclitaxel) x 6 Lipodox x 2 (Bangkok) Plan Lipodox 3 rd course → Private hospital →	1.Rt half colectomy (14 June,2007) 2. Chemotherapy - carboplatin x 2 -> ifosfamide
Outcome	Alive without disease	Lost to follow up	During treatment

Cancer of The Peritoneum 2007. (continued)

Data	Case 7	Case 8	Case 9
Hospital number	3011940	3028193	2629989
Age (yrs)	54	47	47
Marital status	Married	Married	Single
Parity	3-0-0-3	7-0-0-7	-
Presenting symptoms	Pelvic mass	Abdominal distension	Abdominal distension
Stage	IIIC	IIIC	IIIC
Histology	<u>Tumor</u> : Serous cystadenocarcinoma	Poorly differentiated adenoCA <u>Metastasis to</u> both ovaries, tubes,cervix,corpus,omentum,peritoneum,appendix	<u>Peritoneal biopsy</u> : poorly differentiated carcinoma
Treatment	1. Referred case: Explor lap to tumor resection & sigmoid colon resection & end to end anastomosis (4 May,2007) 2. Chemotherapy: PT (carboplatin+ paclitaxel) x 6	1.TAH&BSO&partial omentectomy & appendectomy & ascites collection (2 June,2007) 2. Chemotherapy Cispaltin + 5 FU	1.Laparoscopic examination to pelvic biopsy (Finding: carcinomatosis peritonei) 2. Chemotherapy: PT (carboplatin+ paclitaxel) x →
Outcome	During treatment, await for investigation	Follow-up at medical oncology unit	Progression → palliative treatment

Cancer of The Peritoneum 2007. (continued)

Data	Case 10	Case 11
Hospital number	3031026	3023109
Age (yrs)	43	68
Marital status	Married	Married
Parity	2-0-1-2	6-0-1-2
Presenting symptoms	Abdominal mass	Abdominal distension
Stage	IIC	IV (liver metastasis)
Histology	<u>Tumor</u> : Clear cell carcinoma	-
Treatment	1. 1.TAH&BSO&partial omentectomy & appendectomy & peritoneal washing (24 Oct,2007) 2. Chemotherapy: PT (carboplatin+ paclitaxel)	Supportive & symptomatic treatment
Outcome	During treatment	Lost to follow up

Cancer of Multiple Primaries Gynecologic Organ

Table 43 : Multiple Primary Gynecologic Cancer 2007

Data	Case 1 (Endometrial CA + Cervical cancer)	Case 2 (CA tube + Ca ovary + endometrial CA)	Case 3 (Ca ovary + endometrial CA)
HN	2965259	3005137	2999120
Age (yrs)	55	70	50
Marital status	Married	Married	Married
Parity	2-0-0-2	2-0-0-2	0-0-1-0
Presenting symptoms	Abnormal PAP smear (HSIL)	Abdominal pain & mass	Abdominal pain
Stage	Endometrial CA IB1 + cervical CA IA1	CA ovary IIC + CA tube IC, endometrial CA IB	CA ovary IIIB + endometrial CA IB
Histology	<u>Cervix</u> : squamous cell CA <u>Corpus</u> : endometrioid CA grade 1	<u>Ovary</u> : endometrioid CA grade 2 <u>Corpus</u> : endometrioid CA grade 3 <u>Tube</u> : endometrioid CA grade 2	<u>Ovary</u> : endometrioid CA grade 1 <u>Corpus</u> : endometrioid CA grade 1
Treatment	1. CKC Dec 9,2007 2. TAH+BSO+appendectomy Feb 8,2007	1. Referred: TAH + BSO + partial omentectomy + appendectomy (May 5,2007) 2 Chemotherapy Carboplatin x 6 courses	1. TAH+BSO+ascites collection + omental biopsy 3 Chemotherapy PT x 6 courses
Outcome	Alive without disease	Alive without disease	Alive without disease

Multiple Primary Gynecologic Cancer 2007 (cont')

Case 4 (CA ovary + endometrial CA)	Case 5 (CA ovary + CA tube)	Case 6 (CA ovary +endometrial CA)	Case 7 (CA ovary +endometrial CA)
3028117	3009434	3021594	3047510
55	37	49	57
Married	Married	Single	Married
5-0-0-5	0	0	0
Abnormal uterine bleeding	Abdominal mass + pain	Abdominal pain	Abnormal uterine bleeding
CA ovary IIC + endometrial CA IB	CAovary IV + CA tube IC?	CA ovary IC + endometrial CA IIB	CA ovary IC + endometrial CA IIIA
<u>Ovary:</u> endometrioid CA grade 3 <u>Corpus:</u> endometrioid CA grade 3	<u>Tube:</u> serouscystadeno CA gr 3 <u>Ovary:</u> serouscystadeno CA gr 2	<u>Ovary:</u> endometrioid CA grade 1 <u>Corpus:</u> endometrioid CA grade 1	<u>Ovary:</u> endometrioid CA grade 2 <u>Corpus:</u> clear cell + endometrioid grade 2
<u>1. Referred case</u> Complete surgical staging (August 29 ,2007) <u>2. Chemotherapy</u> PT x 6	<u>1. Referred case</u> BSO+ +appendectomy (June 25,2006) 2. Chemotherapy PT x 3 3. TAH+Partial omentectomy + Paraaortic node sampling + debulking tumor (Sep 19,2007) 4. PT x 3	<u>1. Complete surgical staging</u> (Oct 4,2007) 2. Chemotherapy Carboplatin x 6 3. Pelvic radiation	<u>1. Complete surgical staging</u> (December 25,2007) 2. Chemotherapy - PT
During treatment	Alive without disease	During treatment	During treatment

PT = carboplatin+ paclitaxel

Gestational Trophoblastic Diseases

- Gestational Trophoblastic Tumors
- Molar Pregnancy

TABLE 44 : Gestational Trophoblastic Tumors in 2007.

No	H.N.	Age (yr)	HCG titer	Diagnosis	Class	FIGO stage	Treatment	Result
1	2988941	25	808	CCA	MGTN Lung metas	III	TAH → EMA x 2	Remission
2	2968047	18	>300,000	Persist mole	NMGTT	I	MTX+FA->Act D x 4-> EMA x 4-> EMA x 6	Remission
3	2658065	30	316,040	CCA	NMGTT	I	EMA x 10	Remission
4	3028707	22	212,630	Persist mole	NMGTT	I	MTX + FAx 10	Under treatment
5	2989012	37	15,400	Persist mole	NMGTT	I	MTX + FAx 7	Remission
6	2911373	25	54	Persist mole	NMGTT	I	MTX + FAx 1	Remission
7	2978878	32	12,020	CCA	MGTN Lung metas (wedge resection)	III	EMA x 11	Remission
8	3041684	31	7,701	Persist mole	NMGTT	I	MTX + FAx 1	Lost to follow up
9	3025911	49	4,543	Persist mole	NMGTT	I	MTX + FAx 3-> Act D x 4	Remission
10	3031576	53	199	Persist mole	NMGTT	I	MTX + FAx 4	Remission
11	2970058	21	661	Persist mole	NMGTT	I	MTX + FAx 4-> Act D x 5	Remission
12	2972565	23	813	Persist mole	NMGTT	I	MTX + FAx 1-> Act D x 5	Remission
13	2999865	27	478,497	Invasive mole	MGTN Lung metas	III	TAH → EMA x 9	Remission
14	2965869	50	19,540	Persist Mole (recurrence) Rx	MGTN Lung metas	III	TAH&BSO -> EMA/CE x 17-> 7 mo -> BEP x 4-> refer-> ICE (carboplatin) x 1	Supportive & symptomatic Rx
15	3013994	21	55	Persist mole	NMGTT	I	MTX + FA x 3	Remission
16	2999463	25	37,127	Persist mole	NMGTT	I	MTX + FA x 5	Remission
17	3014289	21	23,247	Persist mole	NMGTT	I	MTX + FA x 1-> Act D x 6	Remission

NMGTN = Nonmetastatic Gestational Trophoblastic Neoplasia
Persist. mole = Persistent mole
MTX + FA = Methotrexate + Folinic Acid
EMA = Etoposide + Methotrexate + Actinomycin D
Rx = Treatment

CCA = Chorio carcinoma
Act D = Actinomycin D
PI = Cisplatin + Ifosfamide
PE = Cisplatin + Etoposide
mo. = months

TABLE 45 : Molar Pregnancy in 2007.

No	Hospital Number	Age (yr)	Gravida	GA (wk)	UT. Size (wk)	HCG titer	Risk	Treatment	Pathology	Result
1	2804127	23	1	16	10	37,120	Low	Dilated & curettage	Partial mole	Remission
2	2953383	31	1	12	6	123,000	High	Dilated & curettage	Partial mole	Remission
3	2791205	18	2	12	12	200,400	High	Dilated & curettage	Complete mole	Refer
4	2957471	20	2	12	14	>500,000	High	Suction & curettage	Complete mole	Lost to follow up
5	2940235	22	1	16	14	10.910	Low	Dilated & curettage	Partial mole	Remission

UT = uterine

yr = years

wk = weeks

SECTION II

- **Medical Personnel and Facilities**
- **Diagnostic Procedures**
and Gynecologic Oncology Operations
- **Publications & Presentations**

Medical Personnel and Facilities

TABLE 46 : Medical Personnel and Facilities
in Division of Gynecologic Oncology, Chiang Mai University

Personnel and Facilities	Number
Medical Doctor	8
General Nurse	28
Practical Nurse	24
Helper	9
Research Nurse	1
Research Assistant	1
Inpatient Bed	62
Outpatient Bed	7
Colposcope	3
Cryosurgery Set	1
Radiosurgery (Surgitron)	2

Funds (กองทุนของหน่วยมะเร็งวิทยา)

1. Gynecologic Cancer Fund (กองทุนมะเร็งทางรีเวช)
2. Cervical Cancer Surgery Fund (กองทุนผ่าตัดมะเร็งปากมดลูก)

1st Year Fellow

- Kannika Paengchit, M.D.
- Kriengkrai Sittidilokratna, M.D.
- Nuttavut Kantathavorn, M.D.

2nd Year Fellow

- Irene Ruengkachorn, M.D.
- Jitima Tiyayon, M.D.

Radiation Oncologists

1. Associate Professor Vicharn Lorvidhaya, M.D.
2. Professor Vimol Sukthomya, M.D.
3. Assistant Professor Anan Tonusin, M.D.
4. Imjai Chitapanarux, M.D.
5. Pimkhuan Kamnerdsupaporn, M.D.
6. Ekkasit Tharavijitkul, M.D.

Gynecologic Pathologists

1. Associate Professor Sumalee Siriaunkgul, M.D.
2. Associate Professor Surapan Khunamornpong, M.D.
3. Associate Professor Jongkolnee Settakorn, M.D.
4. Assistant Professor Kornkanok Sukpan, M.D.

Medical Oncologists

1. Professor Sumitra Thongprasert, M.D.
2. Assistant Professor Chaiyut Charoentum, M.D.
3. Assistant Professor Busyamas Chewaskulyong

Diagnostic Procedures and Operations

TABLE 47 : Diagnostic Procedures and Operations for Cervical Neoplasia.

Procedures & Operations	Number
Colposcopy	519
LEEP	317
Cervical Conization	15
TLH	1
Simple Hysterectomy	47
Radical Parametrectomy & PL	1
Laparoscopic Radical trachelectomy & PL	1
Abandoned Radical Hysterectomy & PL	8
Laparoscopic Radical Hysterectomy & PL	11
Radical Hysterectomy & PL	123
Extended Hysterectomy & PL	18

LEEP = Loop Electrosurgical Excision Procedure

TLH = Total Laparoscopic Hysterectomy

PL = Pelvic Lymphadenectomy

Table 48 : Operations for Ovarian, Corpus, Vagina and Vulvar Cancer and GTT.

Operations	Number
CRS for Ovarian Cancer	81
CRS for Fallopian Tube Cancer	2
CRS for Peritoneal Cancer	6
Surgical Staging for Corpus Cancer	80
Simple hysterectomy for GTT	1
Skinning vulvectomy for Vulva Cancer	1
Wide Local Excision for Vulvar Cancer	1
Radical Hemivulvectomy & BGND for Vulvar Cancer	1
Radical Local Excision & BGND for Vulvar Cancer	2
Radical vulvectomy + BGND for Vulvar Cancer	2
Bilateral Groin Node Dissection for Vulvar Cancer	1

CRS = Cytoreductive Surgery

BGND = Bilateral Groin Node Dissection

GTT = Gestational trophoblastic tumor

**PUBLICATIONS
&
PRESENTATIONS**

1997-2006

THERMAL INJURY IN CERVICAL SPECIMENS OBTAINED FROM LOOP ELECTROSURGICAL EXCISION PROCEDURE (LEEP)

Jatupol Srisomboon*, Sumalee Sirianguk**, Sangwal Ruggpao*, Kasem Ruangrongmorakot*, Prapaporn Suprasert*, Chailert Phongnarisorn*

* Department of Obstetrics & Gynecology,
** Department of Pathology,
Faculty of Medicine, Chiang Mai University

Between August 1, 1995 and November 30, 1997, 178 patients with abnormal cervical cytology were evaluated by colposcopy and underwent loop electrosurgical excision procedures (LEEP) at Maharaj Nakorn Chiang Mai Hospital. The indications for LEEP were unsatisfactory colposcopy (59), pathological evaluation (54), exclusion of invasive cancer (23), discrepancy between cytology and biopsy (14), therapeutic purpose (13), microinvasive carcinoma from biopsy (11), and exclusion of adenocarcinoma (4). Thermal injury at surgical margins could be pathologically assessed in 177 patients. All specimens had some degree of thermal injury including mild degree 91 cases (51.4 %), moderate degree 65 cases (36.7 %), and severe degree 21 cases (11.9 %). Only one specimen was pathologically inevaluable due to severe thermal injury. In conclusion, LEEP provided high rate of thermal injury at surgical margins, but most of the cervical specimens were pathologically evaluable.

Published in : Thai Cancer Journal 1997; 23 : 53-57.

ADENOCARCINOMA OF THE UTERINE CERVIX : A CLINICOPATHOLOGICAL STUDY

Sumalee Siriaungkul*, Sumonmal Maleemonkol**, Surapan Khunamornpong*, Viruch Charoeniam**, Pawares Isariyodom**, Aree Pantusart**

* Department of Pathology,
** Department of Obstetrics and Gynaecology
Faculty of Medicine, Chiang Mai University

Objective : To determine factors which correlated to the risk of recurrence for adenocarcinoma of the uterine cervix treated initially by surgery.

Design : A retrospective study. **Setting:** University Hospital.

Subjects : The clinical information and pathologic specimens of adenocarcinoma of the uterine cervix were retrospectively reviewed for 74 patients who initially had been treated by radical hysterectomy with pelvic lymphadenectomy between 1983 and 1994.

Main outcome measures : Significance of variables in relation to recurrence.

Results : Significant differences in risk of recurrence were observed among the patients with and without lymph node metastasis (50 % vs 15.2 %, $P = 0.03$), and the tumors with increased histologic grade (grade 1 = 12.3 % vs grade 2 + 3 = 41.2 %, $P = 0.01$).

Conclusion : Nodal metastasis and histologic grade were related to recurrence in surgically treated cervical adenocarcinoma.

Published in : Thai Journal of Obstetrics and Gynaecology 1997; 9 : 133-137.

WELL DIFFERENTIATED VILLOGLANDULAR ADENOCARCINOMA OF THE UTERINE CERVIX : A FIRST REPORT OF LYMPH NODE METASTASIS IN TWO OF FOURTEEN CASES.

S.Siriaungkul, S. Maleemonkol,* S. Khunamornpong, V. Charoeniam,* P. Isariyodom,* A. Pantusart*

Department of Pathology,*
Department of Obstetrics and Gynecology
Faculty of Medicine, Chiang Mai University

Background : Well differentiated villoglandular adenocarcinoma is a new subtype of cervical adenocarcinoma. Based on the literature review, neither recurrence nor extrauterine spread has been reported.

Design : The clinicopathological features of well-differentiated villoglandular adenocarcinoma treated by surgery in our institution between 1983-1994 were retrospectively reviewed.

Results : Fourteen cases were included in this study. The patients' age ranged from 22 to 53 (mean, 39.5). Thirteen patients were clinically stage IB and the other was IIA. Two patients had metastatic carcinoma in the pelvic lymph nodes at the time of operation. The tumor invaded deeply to outer third of cervical wall in one case and to middle third with involvement of the entire exocervical mucosa (4 cm. in size) in the other.

Conclusion : Pelvic nodal metastasis could be occurred in the well differentiated villoglandular adenocarcinoma. The depth of invasion is an important factor associated with increased risk of nodal metastasis.

Presented at : Fifth Congress of Asia Pacific Association of Societies of Pathologists & Ninth National Congress of Pathology, December 5-7, 1997 Asia Hotel, Bangkok, Thailand.

THE CLINICAL BENEFIT OF A REPEATED PAPANICOLAOU SMEAR AT THE TIME OF COLPOSCOPY.

Darawadee Ployleumsaeng, Jatupol Srisomboon, Chailert Phongnarisorn, Prapaporn Suprasert

Gynecologic Oncology Unit, Department of Obstetrics and Gynecology, Faculty of Medicine, Chiang Mai University

Abstract : The clinical benefit of a repeated Papanicolaou smear (Pap smear) at the time of colposcopy was determined in 71 patients who had undergone colposcopic examination at the Chiang Mai University Hospital between January 1, 1997 and March 31, 1998. The correlation between the initial Pap smear, the repeated Pap smear and the colposcopically directed biopsy (CDB) results were analyzed. Only 32 patients (45 %) showed an exact correlation between the initial Pap smear and the repeated Pap smear. The sensitivity of the repeated smear was lower than that of the initial smears, which were 76 % and 96 % respectively. The false-negative rate of repeated smear was 23 %. The exact correlation between the first Pap smear and biopsy results did not differ from that of the repeated smear (53.5 % and 50.7 % respectively). Only 2 patients (2.8 %) needed conization when performing a repeated smear. No invasive cancer was missed from skipping the repeated smear. In conclusion, this study demonstrates that repeating of a Pap smear at the time of colposcopy provides a limited clinical benefit. Such procedure can be omitted to save the national budget and reduce workload.

Published in : Chiang Mai Medical Bulletin 1998; 37 (1-2) : 1-5

MALIGNANT LIPID CELL TUMOR OF THE OVARY.

Prapun Jutavijittum, Sumalee Siringkul.

Department of Pathology, Faculty of Medicine, Chiang Mai University

Lipid (steroid) cell tumors of the ovary are one of the rarest forms of ovarian neoplasm, account for less than 0.1 % of all ovarian neoplasms. A 66-year-old Thai woman with pelvic mass was diagnosed as malignant lipid (steroid) cell tumor of the right ovary. The right ovarian mass weighed 850 gm and measured 15x15x10 cm. the tumor composed of two cell types arranged in clusters and groups as organoid pattern intermingled with delicate fibrovascular stroma. One cell type was polygonal, contained eosinophilic and slightly granular cytoplasm. The other cell type was larger, with abundant, vacuolated, often spongy cytoplasm. Foci of necrosis and hemorrhage were observed. The histopathological evidence of metastasis to omentum, wall of sigmoid colon and mesentery of caecum was compatible with malignant lipid cell tumor of the ovary. The patient received 6 courses of chemotherapy and loss to follow-up 7 months after the main operation.

Published in : Thai Journal of Obstetrics and Gynaecology 1998; 10 : 239-242.

OVARIAN MUCINOUS INTESTINAL TUMORS OF LOW MALIGNANT POTENTIAL WITH MICROINVASION: A CLINICOPATHOLOGIC STUDY OF 12 CASES.

Sumalee Siringkul*, Surapan Khunamornpong*, Sumonmal Maleemonkol**, Jatupol Srisomboon**

* Department of Pathology, Faculty of Medicine Chiang Mai University
** Department of Obstetrics and Gynecology, Faculty of Medicine Chiang Mai University.

Objective : To study the clinicopathologic features of the mucinous intestinal tumors of low malignant potential (MILMP) with microinvasion of the ovary.

Design : A descriptive study

Sample : Ovarian MILMP tumors with microinvasion treated at Maharaj Nakorn Chiang Mai Hospital between January 1992 to December 1997 were reviewed.

Result : During six years, there were twelve MILML tumors with microinvasion treated in our hospital. The patients' age ranged from 21-72 years (mean, 44.3) Five of twelve were postmenopausal at the time of surgery. Only two of eleven which had available data were nulliparous. Ten of tumors were classified as stage IA and the other two were IC. The size of tumor ranged from 12-30 cm.(mean 17.1). Seven tumors were right sided, five were on the left. All of them were multilocular cysts with areas of fine honeycombing on the cut surface. Small foci of stromal invasion measured up to 0.3 cm. were identified microscopically. Three of them received adjuvant chemotherapy (two were stage IC and one was IA). Follow up information showed that seven patients were alive without evidence of disease at 7-60 months (mean 19.7), four were lost without evidence of disease at 1-41 months (mean 13.3) and one was lost after surgery.

Conclusion : MILMP tumors with microinvasion are not rare tumors and although the number is small in this study, the prognosis seems to be similar to that of MILMP.

Presented at : XIIIth Annual Scientific Meeting of The Royal Thai College of Obstetricians and Gynaecologists, October 20-22, 1998, Sofitel Raja Hotel, Khon Kaen, Thailand.

MOLAR PREGNANCY IN HILLTRIBE THAI PEOPLE : PROBLEMS AND MANAGEMENT

Srisomboon J, Ployleumsaeng D, Phongnarisorn C, Suprasert P, Puntusart A

Department of Obstetrics & Gynecology,
Faculty of Medicine, Chiang Mai University

Between January 1984 and December 1997, 20 cases of molar pregnancy in hill-tribe Thai people were treated at Department of Obstetrics and Gynecology, Maharaj Nakorn Chiang Mai Hospital. Eight cases (40 %) were below 20 years of age. The mean age was 24.8 years (range 14-50). Eleven cases (55 %) came from Mae Hong Son Province. Twelve cases (60 %) were multiparity. Nineteen cases (95 %) were classified as high risk for malignant sequelae or persistent trophoblastic disease (PTD). Among 6 cases undergone suction curettage without chemoprophylaxis, 2 developed PTD which could be treated until remission. The remaining 4 cases were lost to follow-up after pregnancy termination. Eleven patients were treated by suction curettage with chemoprophylaxis, resulted in remission in 5 cases, in which 2 were cured by continuation of chemotherapy for 3 courses after pregnancy termination. Two patients developing PTD, attained remission following chemotherapy. The remaining 4 patients were lost to follow-up. Three patients underwent abdominal hysterectomy accompanied with chemoprophylaxis. One could achieve remission. The remaining 2 patients were lost to follow-up. Overall, the loss of follow-up rate was 50 percent. Health education policy, especially contraception and antenatal care should be launched to hill-tribe people. Chemoprophylaxis may be considered in hill-tribe patients because of high rate of follow-up loss.

Published in : Bulletin of the Department of Medical Services 1999; 24 : 44-49.

A 14-YEAR RETROSPECTIVE STUDY OF MOLAR PREGNANCY IN MAHARAJ NAKORN CHIANG MAI HOSPITAL : HIGH INCIDENCE OF PERSISTENT DISEASE

Srisomboon J, Ployleumsaeng D, Suprasert P, Phongnarisorn C, Pantusart A.

Department of Obstetrics & Gynecology,
Faculty of Medicine, Chiang Mai University

Objective : To determine the clinical characteristics and outcomes of patients with histologically diagnosed hydatidiform mole.

Design : Retrospective study. **Setting :** Department of Obstetrics and Gynecology, Maharaj Nakorn Chiang Mai Hospital.

Subjects : One hundred and sixty nine patients with histologically diagnosed hydatidiform mole between January 1984 to December 1997

Main outcome measures : Clinical characteristics, remission, and the incidence of persistent gestational trophoblastic disease (PGTD).

Results : The incidence is 1.96:1,000 deliveries. The mean maternal age was 27.8 years (range 14-45). Forty two percent of patients were primigravida. The mean gestational age at diagnosis was 14.4 weeks (range 6-28). The average uterine size at surgical evacuation was 17.4 weeks (range 8-32). The most common presenting symptoms was vaginal bleeding. About 90 % of patients were classified as high risk for PGTD. The incidence of PGTD was 53.3 % in patients undergoing suction curettage without chemoprophylaxis. When separately analyzed, PGTD occurred in 59 % and 16.7 % of high-risk and low-risk patients, respectively. The incidence of PGTD remained high at 43.8 % despite receiving chemoprophylaxis during suction evacuation. PGTD also occurred in 12.5 % of patients undergoing hysterectomy with chemoprophylaxis. Loss of follow-up rate was relatively high at 28 %.

Conclusion : Due to high incidence of PGTD, regular follow-up with sensitive hCG assay must be emphasized to all patients following termination of molar pregnancy.

Presented at : XIIIth Annual Scientific Meeting of The Royal Thai College of Obstetricians and Gynaecologists, October 20-22, 1998, Sofitel Raja Hotel, Khon Kaen, Thailand and IVth International Conference of Nepal Society of Obstetricians & Gynaecologists, February 21-22, 1999, Hotel Himalaya, Kathmandu, Nepal.

Published in: Thai Journal of Obstetrics and Gynaecology 1999; 11:17-22.

MANAGEMENT OF PATIENTS WITH POSITIVE MARGINS AFTER CERVICAL CONIZATION : A REVIEW.

Prapaporn Suprasert, Jatupol Srisomboon.

Division of Gynecologic Oncology, Department of OB&GYN, Faculty of Medicine, Chiang Mai University

Management of patients with positive cone margins is based on the pathology of cone margins, the results of endocervical curettage (ECC) and surgical methods of cervical removal. In case of microinvasive carcinoma (MIC), the risk of residual-invasive cancer is unacceptably high if either cone margins or ECC is positive. A repeated conization or modified radical hysterectomy is recommended depending on the patient's desire of future fertility. For cervical intraepithelial neoplasia (CIN), the risk of residual tumor appears to depend on the method of cervical tissue removal. If positive margins are noted after loop electrosurgical excision procedure (LEEP), further excisional conization is advised to exclude residual disease. Nevertheless, in case of positive margins following cold knife conization (CKC), close follow-up with cervical cytology and colposcopy may be appropriate in patients whose fertility should be retained. For adenocarcinoma in situ (AIS), CKC providing larger cervical specimens is preferred for accurate histological diagnosis. Although the risk of residual disease and recurrence after LEEP is significantly greater than that after CKC, hysterectomy with long term follow-up is recommended in case of positive AIS at surgical margins.

Published in : Thai Journal of Obstetrics and Gynaecology 1999;11 : 53-60.

EXPERIENCE WITH RADICAL HYSTERECTOMY AND PELVIC LYMPHADENECTOMY FOR CERVICAL CANCER WITH NO PERITONIZATION AND NO RETROPERITONEAL DRAINAGE.

Jatupol Srisomboon , Prapaporn Suprasert , Chailert Phongnarisorn

Division of Gynecologic Oncology, Department of OB&GYN, Faculty of Medicine, Chiang Mai University

Objective : To determine postoperative morbidity and lymphocyst formation in cervical cancer patients undergoing radical hysterectomy and pelvic lymphadenectomy (RHPL) without reoperation and retroperitoneal drainage.

Design : Prospective study.

Setting : Division of Gynecologic Oncology, Department of Obstetrics & Gynecology, Faculty of Medicine, Chiang Mai University.

Subjects : Fifty five consecutive patients with stage IA2-IIA cervical cancer underwent RHPL between March, 1997 and November 1998. No patient received preoperative radiation therapy or chemotherapy. A Piver-Rutledge type II-III extended hysterectomy was performed with nonclosure of the pelvic peritoneum. No closed-suction retroperitoneal drainage was utilized after pelvic lymphadenectomy. Sonographic examination was performed at 2, 4 and 6 weeks postoperatively.

Results : The patients were in stage IA2 (1), IB1 (51), and IIA (3) of cervical cancer. The operating time ranged from 2.3 hrs. to 5.2 hrs. (mean = 3.7 hrs). Injury to the bladder occurred in 3 patients. One patient had ureteral injury, another one had rectal injury. The blood transfusion rate was 52.7 %. Number of lymph nodes removed ranged from 10-42 (mean = 26). Postoperative stay ranged from 3-8 days with a mean of 4.3 days. No patient developed bowel obstruction. One patient had febrile morbidity caused by urinary tract infection. Early asymptomatic lymphocysts detected on ultrasound occurred in 2 patients at 6 weeks postoperatively and resolved spontaneously 6 weeks and 8 weeks later. However, one patient complaining of right leg edema at a 6-month follow-up visit was found by CT scan to have a lymphocyst. The lymphocyst and symptom disappeared after fine needle aspiration under ultrasound guidance.

Conclusion : Routine reoperation and retroperitoneal drainage following radical hysterectomy and pelvic lymphadenectomy may be safely abandoned.

* **Published in :** Thai Journal of Obstetrics and Gynaecology 1999; 11 (Suppl.1) : 69-74.

* **Presented at :** XIIIth Annual Scientific Meeting of The Royal Thai College of Obstetricians and Gynaecologists, October 20-22, 1998, Sofitel Raja Hotel, Khon Kaen, Thailand.

SIGNIFICANCE OF SURGICAL MARGIN STATUS IN CERVICAL SPECIMENS OBTAINED FROM LOOP ELECTROSURGICAL EXCISION PROCEDURE (LEEP).

Prapaporn Suprasert ,* Jatupol Srisomboon ,* Sumalee Siriaunkgul ,** Kasem Ruangrongmorakot ,* Chailert Phongnarisorn .*

*Department of OB&GYN,

**Department of Pathology

Faculty of Medicine, Chiang Mai University

Objective : To determine the significance of surgical margin status in cervical specimens obtained from loop electrosurgical excision procedure (LEEP) in patients with cervical neoplasia undergoing such procedure followed by hysterectomy.

Design : Retrospective study.

Setting : Chiang Mai University Hospital.

Method : Between August 1, 1995 and September 30, 1997, ninety-three patients with abnormal Papanicolaou smear underwent colposcopy followed by LEEP. Subsequent hysterectomy was performed within 6 months of the procedure. Medical records were reviewed for patient profiles, the indication of LEEP, pathology of loop specimens including the status of surgical margins and residual disease in subsequent hysterectomy specimens.

Results : Of the 93 patients, 71 were eligible for histologic evaluation of the ectocervical and endocervical margins. The incidence of residual tumor in hysterectomy specimens was highest in patients with positive endocervical margins only (12 of 29 patients or 41.4 %). Of 10 cases with both positive both endocervical and ectocervical margins, 3 (30 %) had residual disease. In 22 cases with both negative endocervical and ectocervical margins, residual tumors were still found in 4 patients (18.2 %). In contrast, 2 out of 10 patients (20 %) with positive ectocervical margins only had residual disease. However, three cases of invasive cancer were missed from such procedure. The first was initially diagnosed as high-grade squamous intraepithelial lesion (HGSIL) with involved endocervical margin. This patient was mistakenly treated with simple hysterectomy. The remaining two patients also had HGSIL but squamo-columnar junction could not be identified due to shallow loop specimens. Both patients underwent radical hysterectomy and inappropriate simple hysterectomy, respectively.

Conclusion : The presence of disease at surgical margins of loop specimens significantly correlates with the residual disease in subsequent hysterectomy specimens. The status of the internal or endocervical margins appears to be more predictive of residual disease on subsequent histologic evaluation when compared with that of the external or ectocervical margins.

Published in : Thai Journal of Obstetrics and Gynaecology 1999; 11 (Suppl 1) : 75-81.

METASTATIC HEPATOCELLULAR CARCINOMA OF THE OVARY.

S. Khunamornpong *, S. Siriaunkgul *, A. Chunduan **

*Department of Pathology, **Department of OB&GYN, Faculty of Medicine, Chiang Mai University

Hepatocellular carcinoma is an uncommon tumor to metastasize to the ovary and may cause a diagnostic problem. To our knowledge, there were six cases of metastatic hepatocellular carcinoma to the ovary reported in the English literature. We, herein, present an additional case in which ovarian lesion was the initial presentation.

A 56-year-old postmenopausal woman presented with pelvic mass for 1 month. Physical examination disclosed mild anemia and lower abdominal mass. The liver was not palpable. Pelvic examination revealed a 15-cm left adnexal mass. An exploratory laparotomy disclosed a left ovarian tumor with 1000 ml. ascitic fluid and carcinomatous peritonei involving the surface of omentum, liver and diaphragm. The tumor extended toward the posterior surface of the uterus and sigmoid colon. The ovary was normal size but contained a few small surface nodules, measuring 3 to 4 mm across. TAH & BSO and sigmoid colectomy were performed.

On microscopic exam, the tumor showed a predominant solid and trabecular patterns with endothelial-bounded vascular spaces. Psudoglandular arrangement was also observed. The neoplastic cells were polygonal-shaped and contained moderate to abundant pale eosinophilic, granular to clear cytoplasm and vesicular nuclei with prominent nucleoli. Evidently observed were intercellular canaliculi, many of which contain bile material that was Hall's stain positive. Mucicarmine stain was uniformly negative. Strong reactivity of the tumor cells for alpha-fetoprotein (AFP) with absence of immunoreactivity for carcinoembryonic antigen (CEA) was noted. Postoperative CT scan revealed multiple confluent masses occupying the left hepatic lobe with several discrete nodules in the right lobe and portal venous thrombosis, consistent with hepatocellular carcinoma. Her serum AFP was 534 ng/ml. The serum was positive for HBsAg, HBeAg, and anti-HBcAb. The patient received supportive treatment with oral megestrol acetate. She died 4 months postoperatively.

Published in : International Journal of Gynecology and Obstetrics
1999; 64 : 189-191.

CLEAR CELL CARCINOMA OF THE OVARY

Sumonmal Manusirivithaya,** Viruch Charoeniam,*
 Pawares Isariyodom,* Jatupol Srisomboon,* Aree Pantusart,*
 Chantawat Sheanakul,** Manit Sripramote,** Surawut Leelahakorn,**
 Thaovilai Thavaramorn,** Siriwan Tangchitkamol**

* Department of Obstetrics and Gynaecology, Faculty of Medicine,
 Chiang Mai University.

** Bangkok Metropolitan Administration Medical College and Vajira
 Hospital.

Objective : To evaluate the clinical characteristics and results of treatment of
 ovarian clear cell carcinoma.

Method : The detailed data on 59 patients with clear cell carcinoma of the ovary
 treated at Maharaj Nakorn Chiang Mai Hospital (between 1984-1996) and Vajira
 Hospital (between 1992-1998) were retrospectively reviewed.

Results : The median age was 46 years. Fifty nine percent were nulliparous.
 The incidence by surgical stage were 33 (55.9 %) stage I (Ia 14, Ic 19), 4 (6.8 %) stage II (IIa 1, IIb 1, IIc 2) and 22 (37.3 %) stage III (IIIa 1, IIIb 4, IIIc 17). The tumor size ranged from 6 to 30 cm. with the median of 12 cm. Bilateral ovarian involvement was found in 3.1 % for stage I and 17.2 % in all stage. Treatment consisted of cytoreductive surgery (with optimal surgery in 81.5 %) followed by chemotherapy in most cases. Considering 37 cases of stage I and II; 3 had tumor progression during treatment, 7 were lost during treatment and 2 of these had tumor progression, 6 had tumor recurrence after complete treatment while 21 were alive without disease (median follow up time 39 months). For 22 cases of stage III; 7 were lost during treatment and all but 1 were known to have tumor progression. Another 13 patients had tumor progression during treatment while only 2 patients (both of stage IIIb) were alive without disease 5 and 10 years after treatment. The 5-year survival was 91.7 % for stage I, 50 % for stage II and 14.4 % for stage III.

Conclusion : Clear cell carcinoma of the ovary was more common presented in early stage and patients in early stage had good prognosis with appropriate treatment while patients in advanced stage had poor prognosis despite aggressive chemotherapy.

Presented at : XIVth Annual Scientific Meeting of The Royal Thai College of Obstetricians and Gynaecologists, October 26-29, 1999, The Royal Golden Jubilee Building, Bangkok, Thailand.

RADICAL PARAMETRECTOMY, UPPER VAGINECTOMY AND PELVIC LYMPHADENECTOMY OF INVASIVE CERVICAL CANCER FOLLOWING SIMPLE HYSTERECTOMY.

Jatupol Srisomboon , Chailert Phongnarisorn , Prapaporn Suprasert ,
 Chalong Cheewagriang-grai , Kittipat Charoenkwan, Sitthicha Siriaree

Division of Gynecologic Oncology, Department of OB & GYN,
 Faculty of Medicine, Chiang Mai University

The standard treatment of early invasive cervical cancer included either radical hysterectomy or radiation therapy. Both treatment modalities have been accepted as equally effective in terms of local control and survival. Although radical hysterectomy and pelvic lymphadenectomy (RHPL) appears to be sufficient for early-stage cervical cancer, some condition arises when the invasive lesion is found incidentally in the specimen obtained from simple hysterectomy. This condition, called cut-through hysterectomy, indicates the need for further therapy either radiation or radical re-operation to eliminate the risk of residual disease in the vaginal cuff, paravaginal tissues, paracervical tissues, and pelvic lymph nodes.

Radical re-operation is an interesting option since ovarian function and vaginal pliability can be preserved. This report presents our experience with radical parametrectomy, upper vaginectomy, and pelvic lymphadenectomy without serious perioperative morbidity in 3 patients undergoing inadvertent simple hysterectomy of invasive cervical cancer. Although the operation seems to be technically more difficult than RHPL, it is an attractive and effective option and can be safely performed in selected patients especially young women with no clinical evidence of residual tumor.

Presented at : XIVth Annual Scientific Meeting of The Royal Thai College of Obstetricians and Gynaecologists, October 26-29, 1999, The Royal Golden Jubilee Building, Bangkok, Thailand.

Published in : Thai J Obstet Gynaecol 2000;12(2):141-4

REASONS FOR IMPROPER SIMPLE HYSTERECTOMY IN PATIENTS WITH INVASIVE CERVICAL CANCER.

Jatupol Srisomboon, Prapaporn Suprasert, Chailert Phongnarisorn,
 Aree Pantusart, Chalong Cheewagriang-grai, Kittipat Charoenkwan,
 Sitthicha Siriaree.

Division of Gynecologic Oncology, Department of OB & GYN,
 Faculty of Medicine, Chiang Mai University

Objective : To determine the reasons for improper simple hysterectomy in the presence of invasive cervical cancer.

Methods : The medical records of 70 patients who had undergone simple hysterectomy in the presence of invasive cervical cancer and were referred to Chiang Mai University Hospital between January 1991 and December 1998 were reviewed.

Results : Approximately half of the patients presented with abnormal vaginal bleeding. Failure to perform a Papanicolaou smear before the operation accounted for 35.4 %. Normal pelvic examination in which no gross invasive tumor was observed accounted for 59 %. The most common indication for inappropriate operation was cervical dysplasia. The reasons for inappropriate simple hysterectomy included lack of preoperative cervical cytology (22.8 %), incomplete evaluation of cervical dysplasia or microinvasion on biopsy (21.4 %), false-negative cervical cytology (18.6 %), failure to perform an endocervical curettage following conization (8.6 %), emergency hysterectomy (8.6 %), failure to perform indicated conization (5.6 %), errors in colposcopic examination (4.3 %), incomplete evaluation of an abnormal cervical cytology (1.4 %), failure to perform endocervical curettage following loop conization (1.4 %), failure to review slide (1.4 %), failure to check the pathology report (1.4 %), failure to biopsy a gross cervical lesion, and skipped lesion in upper part of the endocervix (91.4 %). One case of invasive disease was missed for unknown reason.

Conclusion : Most cases of inappropriate hysterectomy resulted from deviation from guideline for cervical cancer detection. Preoperative Papanicolaou smear and strict adherence to the well-established diagnostic protocol for patients with an abnormal smear are advised to prevent such occurrence.

Presented at : XIVth Annual Scientific Meeting of The Royal Thai College of Obstetricians and Gynaecologists, October 26-29, 1999, The Royal Golden Jubilee Building, Bangkok, Thailand.

Published in : J Obstet Gynaecol Reseach 2000 ; 26(3) : 175-80.

HIGH DOSE RATE AFTERLOADING BRACHYTHERAPY IN CARCINOMA OF THE CERVIX. AN EXPERIENCE OF 1992 PATIENTS.

Lorvidhaya V, Tonusin A, Changwiwit W, Chitapanarux I,
 Srisomboon J, Wanwilairat S, Chawapan N, Sukthomya V

Section of Radiation Oncology, Faculty of Medicine,
 Chiang Mai University

Purpose : To report the result of radiation therapy in carcinoma of the cervix treated by external irradiation and high dose rate intracavitary brachytherapy.

Methods and Materials : This is a retrospective analysis of 2063 patients with histologically proven carcinoma of the cervix treated by external irradiation and high dose rate intracavitary brachytherapy between March 1985 - December 1991. The Kaplan-Meier method was used for survival and disease free survival analysis. Late complications in the bowel and bladder were calculated actuarially.

Results : There were seventy-one patients who did not complete the course of irradiation, so only 1992 patients were retrospectively analyzed for survival. There were 2 patients (0.1 %) in stage IA, 211 (10.2 %) stage IB, 225 (10.9 %) stage IIA, 902 (43.7 %) stage IIB, 14 (0.7 %) stage IIIA, 675 (32.7 %) stage IIIB, 16 (0.8 %) stage IVA, and 16 (0.8 %) in stage IVB. The median follow up time was 96 months. The actuarial 5-year disease free survival rate was 79.5 %, 70.0 %, 59.4 %, 46.1 %, 32.3 %, 7.8 % and 23.1 % for stage IB, IIA, IIB, IIIA, IIIB, IVA and IVB respectively. The actuarial 5-year disease free survival rate for stage IB₁ and IB₂ squamous cell carcinoma was 88.7 % and 67.0 % respectively. The actuarial 5-year overall survival rate was 86.3 %, 81.1 %, 73.0 %, 50.3 %, 47.8 %, 7.8 % and 30.8 % for stage IB, IIA, IIB, IIIA, IIIB, IVA and IVB respectively. Pattern of failure revealed 20.8 % local recurrence, 18.7 % distant recurrence and 4 % in both. The late complication rate grade 3 and 4 (RTOG) for bowel and bladder combined was 7.0 % with 1.9 % grade 4.

Conclusions : The result in high dose rate brachytherapy used in this series, produced pelvic control and survival rates comparable to other LDR series.

Published in : International Journal of Radiation Oncology, Biology & Physics 2000;46:1185-91

UPDATE RESULT ON GEMCITABINE AND CISPLATIN IN PATIENTS WITH METASTATIC CERVICAL CANCER

V. Lorvidhaya, I. Chitapanarux, P. Kammerdsupaphon.

Division of Therapeutic Radiology and Oncology, Faculty of Medicine, Chiang Mai University

Purpose : To determine the therapeutic efficacy of cisplatin plus gemcitabine in the treatment of patients with metastatic cervical cancer.

Methods and Materials : A total of 51 patients were enrolled in this study. The median age was 46 years (34-67). Thirty-five patients were treated to the pelvis by radical radiotherapy, one patient was treated by surgery and one by surgery and postoperative RT previously. There were 14 patients with stage IVB cervical cancer who were previously untreated. The sites of metastases were 10 in the lungs, 5 in the supraclavicular nodes, 9 in the paraaortic nodes, 4 in the liver, 3 in the inguinal nodes, 5 in both supraclavicular nodes and lungs, 9 in both supraclavicular and paraaortic nodes, 1 in both supraclavicular and inguinal nodes, 1 in both lung and inguinal nodes, 2 in both lung and paraaortic nodes and 2 in both liver and lungs. Fine needle biopsies were done in all metastatic sites except for multiple lung, and multiple liver metastases. Cisplatin was administered as IV infusion on day 1 (70 mg/m²). Gemcitabine was administered as an IV infusion over 30 minutes on day 1 and 8 (1250 mg/m²) every 21 day cycle.

Results : Thirty-seven patients were available for evaluation : 3/37 (8.1%) achieved a complete response, 26/37(70.3%) achieved a partial response, 4/37 (10.8%) had stable disease, and 4 (10.8%) had progressive disease. Myelosuppression was the major toxicity. Grade 3 or 4 anemia and granulocytopenia occurred at a frequency of 25.5% and 29.4%. high grade thrombocytopenia was found in 3.8%. there were 3 (5.8%) patients who developed grade 4 neutropenia and fever. No other major side effects were found other than alopecia and usual gastrointestinal toxicities such as anorexia, nausea and vomiting.

Conclusion : In this study of patients with metastatic cervical cancer, the combination of cisplatin and gemcitabine induced a high response rate. These results require confirmation in larger number of patients.

Presented at : Lilly Oncology Conference, Update in Cervical Cancer Management, November 17-19, 2000, Kunming, People's Republic of China.

A PROSPECTIVE RANDOMIZED STUDY COMPARING RETROPERITONEAL DRAINAGE WITH NO DRAINAGE AND NO PERITONIZATION FOLLOWING RADICAL HYSTERECTOMY AND PELVIC LYMPHADENECTOMY (RHPL) FOR INVASIVE CERVICAL CANCER (ICC).

Jatapol Srisomboon, Prapaporn Suprasert, Chailert Phongnarisorn, Chalong Cheewagriang-grai, Sitthicha Siriaree, Kittipat Charoenkwan, Aree Pantusart

Division of Gynecologic Oncology, Department of OB & GYN, Faculty of Medicine, Chiang Mai University

Objective : To evaluate the postoperative morbidity and lymphocyst formation in ICC patients undergoing RHPL with no drainage and no peritonization compared with retroperitoneal drainage and peritonization.

Methods : Between July 1999 and May 2000, 100 patients with stage IA-IIA cervical cancer undergoing RHPL were randomized to receive either no drainage and no peritonization (Group A=48 cases) or retroperitoneal drainage and peritonization (Group B = 52 cases). Perioperative data and morbidity were recorded. Transabdominal and transvaginal sonography were performed at 1,2, and 3 months postoperatively to detect lymphocyst formation.

Results : Both groups were similar regarding age, size and gross appearance of tumor, tumor histology, and stage. There was no difference between groups in respect of operative time, need for blood transfusion, intraoperative complications, number of nodes removed, nodal metastases, and need for adjuvant radiation and chemotherapy. Asymptomatic lymphocysts were sonographically detected at 1,2 and 3 months postoperatively in 3 (6.8%), 2(4.6%), and 3 (7.7%) of 44,43, and 39 patients respectively in Group A, whereas none was found in Group B (P=0.2). no significant difference was found in term of postoperative morbidity in the two groups.

Conclusion : Routine retroperitoneal drainage and peritonization after radical hysterectomy and pelvic lymphadenectomy for invasive cervical cancer can be safely omitted.

Presented at : XVth Annual Scientific Meeting of The Royal Thai College of OB&GYN, October 17-20, 2000, Dusit Island Resort Hotel, Chiang Rai, Thailand.

Published in : J Obstet Gynaecol Res 2002;28:149-53.

ATYPICAL SQUAMOUS CELLS OF UNDETERMINED SIGNIFICANCE : CHIANG MAI EXPERIENCE.

Charin Ya-In, Benjaporn Chaiwun, Jonkolnee Settakorn , Samreung Rangaeng .

Department of Pathology, Faculty of Medicine, Chiang Mai University

Cervicovaginal smears have been widely accepted as the most effective screening method for carcinoma of cervix. A new reporting system of cervicovaginal smear, so called "The Bethesda System" was established since 1988. The term "Atypical squamous cells of undetermined significance or ASCUS" was introduced in this reporting system for those squamous cells that were abnormal but did not fulfil criteria for diagnosis of dysplasia or squamous intraepithelial lesion.

The Bethesda System had been used in our institution since mid 1995. Following some adjustment, we would like to evaluate whether our performance was similar to those published in the literature and also to appraise outcome of ASCUS after follow-up

This brief communication includes our preliminary data concerning ASCUS in our experience from January to October 2000. Out of 12,879 cases screened during this period, 146 cases (1.13%) of ASCUS were diagnosed while 986 cases (7.66%) and 601 cases (4.67%) of high grade squamous intraepithelial lesion (HSIL) and low grade squamous intraepithelial lesion (LSIL) were encountered, respectively. Among 146 ASCUS cases, 79 (54.12%) of HSIL, 8 cases (10.13%) of LSIL and 6 cases (7.59%) of frank invasive squamous cell carcinoma. A single case had inadequate tissue for evaluation.

We conclude that our performances are similar to those reported in the literature. The ASCUS : SIL ratio in only 0.08 that falls in an acceptable range (less than 3). Approximately one-thirds (31.0% of our ASCUS cases have precancerous or cancerous histologic follow-up. The results resemble some reported series in the western country.

Presented at : The Seventh Thai-Japanese Workshop in Diagnosis Cytopathology and The First Annual Congress of Thai Society of Cytology, January 10-12, 2001, Phuket Merlin Hotel, Phuket, Thailand.

MALIGNANT OVARIAN NEOPLASMS: HISTOLOGIC SUBTYPES OF 314 CASES TREATED AT THE UNIVERSITY HOSPITAL OF NORTHERN THAILAND.

Siriaungkul S.¹ Khunamornpong S.¹ Srisomboon J.² Wisedmongkol W.¹

Department of Pathology¹ Department of Obstetrics and Gynecology² Faculty of Medicine, Chiang Mai University

To investigate the relative frequency of each histologic type of malignant ovarian neoplasms at Maharaj Nakorn Chiang Mai Hospital. All cases of primary ovarian malignant tumors treated in our hospital during 6-year-period (1992-1997) were evaluated for the histologic types. Of 314 malignant tumors, 248 (79.0%) were of epithelial-stromal group, 14 (4.4%) were sex cord-stromal tumors, and 50 (16.0%) were germ cell tumors. Among 138 invasive carcinomas, the major types included endometrioid (29.0%), serous (27.5%), clear cell (19.6%), and mucinous (11.6%) adenocarcinoma. Of 110 LMP tumors, 75.45 were mucinous; 14.5%, serous; 7.3%, endometrioid; and 2.7%, mixed LMP tumors. Microinvasion was also observed in 11.8% of LMP tumors. The ratios of serous to mucinous type were 2.4:1 for invasive carcinoma and 0.2:1 for LMP tumors. Of malignant germ cell tumors, dysgerminoma, yolk sac tumor, and immature teratoma were the major types with similar relative frequency. Compared to the data in the literature, less relative frequency of serous adenocarcinoma, and higher relative frequency of mucinous LMP tumors and malignant germ cell tumors was observed in our hospital than in most Western countries.

Presented at : XXIII International congress of the international academy of pathology and 14th world congress of academic and environmental pathology 15-20 October, 2000 Nagoya, Japan.

SECOND LOOK OPERATION FOR OVARIAN CANCER

Sumonmal Manusirivithaya* Siriwan Tangjitkamol*
 Manit Sripramodya* Surawute Leelahakorn*
 Taovalai Thavaramara* Chantawat Sheanakul*
 Viruch Charoeniam** Pawares Isariyodom**
 Jatupol Srisomboon*** Aree Pantusart***

Department of Obstetrics and Gynecology
 * Bangkok Metropolitan Administration Medical College and Vajira Hospital
 ** Kasemradth Hospital, Chiang Mai
 *** Faculty of Medicine, Chiang Mai University

Objective : To determine the clinical outcome of ovarian cancer patients who underwent second look operation.

Methods : Medical records and follow up information of 130 ovarian cancer patients who underwent second look operation between 1984-1996 at Maharaj Nakorn Chiang Mai Hospital and between 1993-2000 at Vajira Hospital were retrospectively reviewed.

Results : The median age of the patients was 45.5 years. Seventy-six percent had common epithelial ovarian cancer, while 21.5% and 2.3% had germ cell and sex cord stromal tumor. The stage distribution were as followed: stage I 46.2%, stage II 11.5%, stage III 38.5% and stage IV 3.8%. Second look operation was reported as negative in 100 cases (76.9%). With the median follow up of 7.2 years, 19.0% of these patients had tumor recurrence. The median time from negative second look operation to recurrence was 1 year (1 month to 5.3 years). Of these recurrent cases, only 5.3% were free of disease at last visit of ≥ 2 years after recurrence. Regarding those 30 cases (23.1%) who had positive second look operation; 23.3% had only cytologic and/or microscopic positive and 16.7% were free of disease at last visit for more than 5 years after second look operations.

Conclusion : The benefit of second look operation is controversial since recurrence can occur in those with negative second look; however, some patients with positive second look will be treated earlier and had long survival.

Presented at : 16th Annual Scientific Meeting of The Royal Thai College of OB&GYN, The Royal Golden Jubilee Building, Bangkok, Thailand. October 16-19, 2001.

A PROSPECTIVE RANDOMIZED STUDY COMPARING VOIDING TIME BETWEEN INTERMITTENT SELF-CATHETERIZATION AND SUPRAPUBIC CYSTOSTOMY FOLLOWING RADICAL HYSTERECTOMY AND PELVIC LYMPHADENECTOMY FOR CERVICAL CANCER

Prapaporn Suprasert, Jatupol Srisomboon, Chailert Phongnarisorn.

Department of OB & GYN, Faculty of Medicine, Chiang Mai University

Objective : To compare the voiding time between intermittent self-catheterization (ISC) and suprapubic cystostomy (SPC) in cervical cancer patients undergoing radical hysterectomy and pelvic lymphadenectomy

Design : Prospective randomized study

Setting : Division of Gynecologic Oncology, Department Obstetrics & Gynecology, Faculty of Medicine, Chiang Mai University.

Methods : Between September 1998 and June 1999, 71 patients with stage IB-IIA cervical cancer undergoing RHPL were prospective randomized to receive either ISC (38 cases) or SPC (33 cases) for managing post operative bladder dysfunction. The intermittent self-catheterization was initiated on 7th post operative day every 4 hours after voiding in the ISC group, whereas in the SPC group, the catheter was clamped on 7th post operative day. The duration for clamping in the SPC group was 4 hours and was released after self voiding to check the residual urine. Both ISC and SPC were discontinued when post voiding residual urine was less than 75 ml for 2 consecutive voids. Patients in the SPC group was switched to perform ISC if they could not void normally at 30 days postoperatively. The voiding time, the frequency of urinary tract infection, perioperative data and morbidity were recorded.

Results : Both groups were similar regarding age, BMI, tumor histology, stage, size and gross appearance of the tumors. There was no difference between groups in respect of operative time, need for blood transfusion, peri-operative morbidity, need for adjuvant chemoradiation and the frequency of urinary tract infection. The mean operative time of the SPC group was significantly longer than the ISC group (3.8 vs 3.5 hr, $P=0.05$). Twenty-five (65.8%) patients in the ISC group and 23 (69.7%) in the SPC group had voiding time less than 30 days. Mean voiding time in the ISC group was significantly less than that in the SPC group (13.1 days vs 17.3 days respectively, $P=0.02$).

Conclusion : Patients performing ISC after RHPL for invasive cervical cancer can resume bladder function earlier than those undergoing SPC without significant difference in the frequency of urinary tract infection and perioperative morbidity.

Presented at : XVIth Annual Scientific Meeting, The Royal Thai College of OB&GYN, The Royal Golden Jubilee Building, Bangkok, Thailand. 19 October, 2001

Published in : Thai J Obstet Gynaecol 2002;14:73-9

HUMAN PAPILLOMA VIRUS DETECTION AND EXPRESSION OF HPV 16 AND 18 E6/E7 mRNA IN CERVICAL CANCER CELLS

Pranee leechanachai, Chatsri Kuansuwan, Sungwal Rugpao*,
 Jatupol Srisomboon*, Vatcharin Suntornlimsiri*,
 Nipapan Komsattum*, Wasun Chantratitta, Wasna Sirirungsri

Department of Clinical Microbiology, Faculty of Associated Medical Sciences,

*Department of Obstetrics and Gynecology, Faculty of Medicine, Chiang Mai University.

Several types of human papilloma virus (HPV) have been implicated in the development of neoplastic lesions of the uterine cervix. HPV type 16 and 18 are the most frequently found especially in cervical cancer cells and are designated as a high-risk types. The viral transforming function are located in E6/E7 open reading frames. The expression of those genes are important for malignant transformation. In this study, we used the polymerase chain reaction based restriction fragment length polymorphism (PCR-RFLP) for detection and typing of the HPV in preinvasive and invasive cervical carcinomas. The HPV 16 and 18 containing cervical cancer cells were further analysed for the patterns of E6/E7 transcription. The relative amount between E6/E7 mRNA subsets was determined by using ABI 310 Genetic Analyzer (Perkin Elmer, U.S.A). In preinvasive stage, CIN1, CIN2, CIN3 and CIS, HPV DNA was detected in 9 out of 50 samples (18%). In this group, HPV 16, 18, 52 and 59 were detected in 2, 2, 4 and 1 sample (s) respectively. For the invasive stage, 32 out of 36 samples (88.9%) were containing HPV DNA. HPV 16 was predominately detected in 15 samples (46.9%) while HPV 18, 35, 52, 6 and 11 were detected in 4, 7, 2, 2 and 1 sample (s) respectively. At least two subsets of mRNA, E6*I and E6*II were detected in all HPV 16/18 containing cervical cancer cells. However, the E6*I mRNA that function in producing E7 protein was predominately expressed. The ratio between E6*I and E6*II was vary from 1 to 2. This study suggested at least HPV especially the high-risk type 16 and 18 together with their E6/E7 oncogene expression were important in development and maintaining malignant phenotype of cervical cancer.

Presented at : 5th Asia-Pacific Congress of Medical Virology at Denpasar-Bali, Indonesia, June 26-28, 2000

RANDOMIZED TRIAL OF PACLITAXEL PLUS PARAPLATIN VERSUS CYCLOPHOSPHAMIDE PLUS PARAPLATIN IN THE TREATMENT OF ADVANCED EPITHELIAL OVARIAN CANCER

Chaiyod Thirapakawong, Somchaya Neungton, Sumrit Senapad,*
 Pornchai Mekariya, Kasem Vichaiathum,** Jatupol Srisomboon.***

* Siriraj Hosital, Mahidol University, Bangkok 10700

** Bhumibol Adulyadej Hospital, Bangkok 10210

*** Maharaj Nakorn Chiang Mai Hospital, Chiang Mai University

Objective : To compare the efficacy and toxicity of paclitaxel plus paraplatin with paraplatin plus cyclophosphamide in the treatment of the advanced epithelial ovarian cancer.

Design : Randomized trial.

Setting : Multicenters.

Subjects : The valuable 95 patients of the stage III-IV epithelial ovarian cancer from 3 centers were recruited between December 1995-April 1998.

Main outcome measure : Response rate, drug toxicity, recurrent rate and survival

Results : The paclitaxel 150 mg/m² plus paraplatin AUC 6 and cyclophosphamide 700 mg/m² plus paraplatin AUC 6 achieved similar results in clinical complete response 63.8% and 66.7%, pathological complete response 76.0% and 75.0%, recurrence after pathological complete response 36.8% and 55.5% respectively ($P>0.05$). The median disease free survival in the pathological complete response group of paclitaxel plus paraplatin cannot be estimated, whereas 11.0 months of paraplatin plus cyclophosphamide. The mean disease free survival of two groups fell in 39.8 and 27.1 months ($p=0.07$). The overall mean survival achieved 36.1 and 35.2 months ($p>0.05$). The toxicity was similar in both arms, except myalgia/arthralgia and skin rash occurrence in paclitaxel plus paraplatin arm.

Conclusion : Paclitaxel plus paraplatin comparing with cyclophosphamide plus paraplatin in the treatment of advanced epithelial ovarian cancer showed no significant difference in complete response, recurrence, disease free survival and median survival with the acceptable toxicity.

Key word : paclitaxel plus paraplatin, cyclophosphamide plus paraplatin, epithelial ovarian cancer

Published in : Thai J Obstet Gynaecol 2000;12:295-302

COMPARATIVE STUDY OF BULKY STAGE IB AND IIA CERVICAL CANCER PATIENTS TREATED BY RADICAL HYSTERECTOMY WITH AND WITHOUT NEOADJUVANT CHEMOTHERAPY: LONG TERM FOLLOW-UP

Sumonmal Manusirivithaya* Pawares Isariyodom**
Viruch Chareoniam** Jatupol Srisomboon *** Aree Pantusart ***

Department of Obstetrics and Gynecology
* Bangkok Metropolitan Medical College and Vajira Hospital
** Kasemradht Hospital Chiang Mai
*** Faculty of Medicine, Chiang Mai University

We retrospectively reviewed 190 patients with bulky (>3cm.) stage IB and IIA cervical cancer patients undergoing radical hysterectomy between 1991 and 1994 at Maharaj Nakorn Chiang Mai Hospital to determine whether neoadjuvant chemotherapy (NAC) with MVAC (Methotrexate, Vinblastine, Adriamycin, Cisplatin) can improve survival in bulky stage IB and IIA cervical cancer patients treated by radical hysterectomy. There were 42 patients treated with preoperative NAC (MVAC 1-3 courses) and 148 patients treated by primary surgery (PS). In the NAC group, the overall response rate from MVAC was 88.1% with 31.0% clinical complete response and 7.1% pathological complete response. Pelvic lymph node metastasis was not significantly different between the NAC group (16.7%) and the PS group (18.2%). At a median follow up of 64.5 months, 19.0% in the NAC group and 18.2% in the PS group had tumor recurrence. The 5-year progression free and overall survival were 80.8% and 92.0% respectively for the NAC group which were not significantly different from those of 80.2% and 92.9% respectively in the PS group. In conclusion, although NAC can decrease the tumor size and produce high response rate, it does not improve survival in bulky stage IB and IIA cervical cancer patients.

Published in : J Med Assoc Thai 2001;84:1550-7

ENDOMETRIAL CANCER DIAGNOSED IN PATIENTS UNDERGOING HYSTERECTOMY FOR BENIGN GYNECOLOGIC CONDITIONS

Jatupol Srisomboon , Chailert Phongnarisorn , Prapaporn Suprasert

Division of Gynecologic Oncology, Department of OB & GYN,
Faculty of Medicine, Chiang Mai University

Objective: To analyze the events preceding incomplete surgical staging procedures in endometrial cancer patients.

Design : Retrospective descriptive study

Setting : Department of Obstetrics & Gynecology, Faculty of Medicine, Chiang Mai University

Subjects : Thirty patients who were unexpectedly found to have endometrial cancer after undergoing hysterectomy for benign conditions and were referred to Chiang Mai University Hospital between January 1993 to December 1999.

Results : Nineteen patients (63%) presented with abnormal uterine bleeding, but only 9 had undergone fractional uterine curettage. The pre-operative pathological reports in 8 patients were endometrial hyperplasia⁽⁵⁾, proliferative endometrium⁽¹⁾, chronic endocervicitis⁽¹⁾, and endometrial polyps⁽¹⁾. The remaining one patient underwent an emergency hysterectomy for severe uterine bleeding following fractional curettage. Eleven patients presented with symptoms other than bleeding, i.e. pelvic mass⁽⁶⁾, pelvic pain⁽³⁾, vaginal discharge⁽¹⁾, and uterine prolapse⁽¹⁾. Ten of the 30 operations were performed for myoma uteri. The preoperative diagnoses of the remaining 20 patients were abnormal uterine bleeding⁽⁵⁾, endometrial hyperplasia⁽⁵⁾, ovarian tumor⁽⁴⁾, pelvic mass⁽³⁾, intractable vaginal discharge⁽¹⁾, uterine prolapse⁽¹⁾, and endometrial polyps⁽¹⁾. Subsequent pathologic evaluation revealed gross appearance of endometrial cancer in the hysterectomy specimens of these patients.

Conclusion : Endometrial cancer might be detected in women undergoing hysterectomy for benign conditions. This situation is best prevented by careful evaluation of patients with abnormal uterine bleeding before definitive surgery. During curettage, the entire endometrium must be removed for accurate pathologic diagnosis. Routine intra-operative opening of the hysterectomy specimen is advised to detect any evidence of endometrial cancer.

Key words : Endometrial cancer, Hysterectomy, Misdiagnosis

Published in : Thai J Obstet Gynaecol 2001;13:29-32

PHASE II TRIAL OF DOCETAXEL AND CARBOPLATIN IN CISPLATIN-RECURRENT ADVANCED OVARIAN CANCER : A PRELIMINARY REPORT

Prapaporn Suprasert, Jatupol Srisomboon , Chailert Phongnarisorn,
Chalong Cheewakriangkrai , Sithicha Siriaree, Sumitra Thongprasert *

Department of OB&GYN, Faculty of Medicine, Chiang Mai University
* Department of Medicine, Chiang Mai University

Background : Patients with recurrent ovarian cancer have a poor prognosis with low response rates and short survival after salvage chemotherapy. In view of encouraging antitumor activity of docetaxel in the preclinical and phase I studies and the synergistic antitumor effect of this agent when combined with carboplatin, we accordingly, conducted a phase II trial of this regimen in patients with cisplatin-recurrent advanced ovarian cancer .

Objective : To evaluate the response rate and toxicity of docetaxel and carboplatin in patients with cisplatin-recurrent advanced epithelial ovarian cancer

Methods : Sixteen patients with cisplatin – recurrent EOC were recruited between October 1999- October 2000. Eligibility criteria included measurable > 2 cm. in diameter EOC. Age > 18 years old, an ECOG performance status of 0-2, adequate hepatic, renal and bone marrow function, and written informed consent. Docetaxel 75 mg/m² as 1 hour IV infusion and carboplatin AUC 5 were given on day 1 every 3 weeks for a maximum 6 cycles. All patients were evaluated for response and WHO grading toxicity.

Results: Eleven patients had recurrence more than 6 months (range 8-24 months) after cisplatin. One patient had disease progression after 2 cycles of cisplatin and 4 had recurrence within 6 months after receiving cisplatin. Four patients were withdrawn from the study because of infusion reaction (1 case), refusal of chemotherapy due to febrile neutropenia (2), and loss to follow up after 4 cycles (1 case). Twelve patients were assessable for response. Five patients had progression disease. The overall response rate was 5 of 12 (41.6%) with 1 CR and 4 PR. The median time to progression was 7 months (2-14 months). Of the 65 cycles, grade 3 or 4 leukopenia and neutropenia occurred in 64.6% and 43.8%, respectively. Febrile neutropenia was found in 23% but only 4.6% required GCSF. Other toxicity was minimal.

Conclusion: The combination of docetaxel and carboplatin appear to have moderate efficacy and manageable toxicity in patient with cisplatin-recurrent epithelial ovarian cancer.

Presented at : 6th Annual Meeting of The Thai Gynecologic Oncology Group.
Felix River Kwai Resort, Kanchanaburi, Thailand, August 11-13,2001

A PROSPECTIVE PHASE II STUDY OF GEMCITABINE PLUS CISPLATIN AS FIRST-LINE CHEMOTHERAPY IN ADVANCED EPITHELIAL OVARIAN AND FALLOPIAN TUBE CANCER: A PRELIMINARY REPORT.

Prapaporn Suprasert, Jatupol Srisomboon, Chailert Phongnarisorn,
Chalong Cheewakriangkrai, Sithicha Siriaree

Department of OB&GYN, Faculty of Medicine, Chiang Mai University

Purpose: We assessed the therapeutic efficacy and toxicity of gemcitabine plus cisplatin (PG) in the treatment of patients with advanced ovarian and fallopian tube cancer (FIGO stage III-IV) who did not receive any previous chemotherapy, radiotherapy or hormonal therapy.

Patients and methods : Eligibility criteria included chemo-naïve patients with advanced epithelial ovarian or fallopian tube cancer, age between 18 and 70 years, an ECOG performance status of 0-2; adequate hepatic, renal and marrow function; and written inform consent. Gemcitabine 1250 mg/m² followed by cisplatin 75 mg/m² were given on day 1 and gemcitabine 1250 mg/m² alone was given on day 8, every 3 weeks for 6 cycles. The patients were evaluated for response (via CT scan and CA125) and toxicity (WHO grading).

Results: Nine patients with ovarian cancer (5 stage IV, 3 stage IIIc and 1 stage IIb) and one with fallopian tube cancer (stage IIIc) were recruited between May 2000 and August 2001. All received chemotherapy after cytoreductive surgery except for 1 patient with fallopian tube cancer who underwent interval debulking surgery after receiving 3 cycles of chemotherapy. Median age was 54 years with a range from 40 to 66 years. Four cases (40%) completed 6 cycles of chemotherapy. Treatment was prematurely stopped after receiving only 1 cycle due to grade 3 mucositis in 2 cases, progression of disease in 1 case and loss to follow up in the other one. Two patients received only 3 and 2 cycles because of grade IV diarrhea and progression of disease respectively. Gemcitabine were administered over 3 hours IV infusion in 25 cycles and within 30 minutes in 8 cycles. Grade 3 and 4 toxicity of anemia (12%), leucopenia (44%), neutropenia (92%), thrombocytopenia (60%), mucositis (8%) and diarrhea (4%) were found in 25 cycles of 3 hours IV infusion of gemcitabine. While only 1 in 8 cycles of 30 minute IV infusion of gemcitabine had grade 3 neutropenia. Of the 4 patients who completed 6 cycles of chemotherapy, the response rate was 75% (2 had complete response, 1 had a partial response, and 1 had progressive disease). Two patients developed progression of disease after receiving 1 and 2 cycles of chemotherapy, respectively.

Conclusion : Gemcitabine plus cisplatin is effective in advanced epithelial ovarian and fallopian tube cancer. Short duration of gemcitabine infusion is recommended to reduce toxicity. The study is ongoing.

Presented at : Second Lilly Oncology Weekend Program: Oncology Thailand Meet China. Shanghai Cancer Hospital. Shanghai, China, 3 November, 2001

ETOPOSIDE, METHOTREXATE, AND ACTINOMYCIN D (EMA) REGIMEN IN MODERATE & HIGH RISK GESTATIONAL TROPHOBLASTIC TUMORS (GTT)

Prapaporn Suprasert, Jatupol Srisomboon, Chailert Phongnarisorn, Sitthicha Siriaree, Chalong Cheewakriangkrai

Division of Gynecologic Oncology, Department of OB & GYN, Faculty of Medicine, Chiang Mai University

Objective: To evaluate the efficacy and safety of etoposide, methotrexate and actinomycin D (EMA) as primary and secondary therapy for moderate and high-risk gestational trophoblastic tumors (GTT).

Methods: In a retrospective study, the medical records of all patients with moderate and high-risk metastatic GTT according to WHO scoring system receiving primary or secondary EMA regimen were reviewed. Hematologic toxicity was graded using WHO criteria.

Results: From December 1997 through May 2001, Six high-risk and two moderate-risk GTT patients received EMA regimen as the primary treatment. One high-risk GTT patient received EMA regimen as secondary therapy after failure to single agent methotrexate from the primary hospital. The median age was 27 years (range 15-55). The regimen was given every 14-21 days. Five cases showed multiple metastatic sites. One case had brain metastasis and the other one had ureter involvement. Six patients (66%) had histologically confirmed diagnosis of choriocarcinoma. Two patients was lost to treatment after three cycle of EMA and all showed the regression of B-hCG. Seven cases received completed treatment and all (100%) achieved complete remission. Only 34 of 50 cycle had the data of hematotoxicity. Grade 3 and 4 neutropenia was found in 10 cycles (29.4%) of EMA regimen. One case had grade 4 anemia which may result from severe hematuria after the first cycle of chemotherapy. Neither grade 3 and 4 leukopenia nor thrombocytopenia was found. There was no other severe toxicity.

Conclusion: EMA regimen was effective in inducing remission of moderate and high risk GTT and the main toxicity is neutropenia which is manageable.

Presented at : 6th National Cancer Conference. Le Royal Meridian Hotel, Bangkok, Thailand. 3-4 December, 2001.

WELL-DIFFERENTIATED VILLOGLANDULAR ADENOCARCINOMA OF THE UTERINE CERVIX: A REPORT OF 15 CASES INCLUDING TWO WITH LYMPH NODE METASTASIS

Surapan Khunamornpong *, Sumalee Siriaunkgul *, Sumonmal Maleemonkol **, Aree Pantusart **

* Department of Pathology

**Department of OB& GYN, Faculty of Medicine, Chiang Mai University

Well-differentiated villoglandular adenocarcinoma is a recently described subtype of cervical adenocarcinoma. The tumor of this type is reported to have distinct clinicopathologic features and excellent prognosis. However, lymph node metastases of this tumor have been described in few reports. Fifteen cases of well-differentiated villoglandular adenocarcinoma treated at Maharaj Nakorn Chiang Mai Hospital were retrospectively reviewed for both clinical and histopathological features. All patients underwent radical hysterectomy with pelvic lymphadenectomy. In the cases with lymph node metastasis, adjuvant radiation therapy was also given. The patients ranged in age from 22 to 53 years (mean, 39.3). Fourteen patients were FIGO stage IB and one was stage IIA. All patients had exophytic friable cervical masses. Tumor size known in 14 cases ranged from 1.5 to 4 cm (mean, 2.3). Eleven tumors (73.3%) were confined to the inner third of the cervical stroma with 9 of these (60%) showing only superficial invasion (depth \leq 3 mm). The tumors invaded deeply to the middle third in 3 cases (20.0%), and to the outer third in one (6.7%). Lymphatic invasion was observed in 3 cases, two of them had pelvic lymph node metastasis. Both patients had tumors involving deeper than the inner third of the cervical wall. The follow-up duration ranged from 21 to 144 months (mean, 67.5). Four of thirteen cases without nodal metastasis were lost to follow-up 36 to 59 months after surgery. All patients showed no evidence of disease at the last visit. Presence of lymphatic invasion and deep stromal involvement appeared to be the risk factors for lymph node metastasis of well-differentiated villoglandular adenocarcinoma.

Presented at: XXIII International congress of the international academy of pathology and 14th world congress of academic and environment pathology 15-20 October, 2000 Nagoya, Japan.

Published in: J Med Assoc Thai 2001; 84: 882-888.

LOOP ELECTROSURGICAL EXCISION PROCEDURE (LEEP) AT MAHARAJ NAKORN CHIANG MAI HOSPITAL : PROBLEMS IN PATHOLOGIC EVALUATION

Surapan Khunamornpong *, Kasem Raungrongmorakot, **, Sumalee Siriaunkgul *

* Department of Pathology

** Department of OB&GYN

Faculty of Medicine, Chiang Mai University

Background : Loop electrosurgical excision procedure (LEEP) is widely used in diagnosis and management of cervical lesions. Difficulties in histopathologic evaluation of LEEP specimens, particularly for the margin status, have been reported to be a significant disadvantage of the procedure.

Method : The histologic slides of the specimens from 163 patients who underwent LEEP at Maharaj Nakorn Chiang Mai Hospital from August 1995 to November 1997 were retrospectively reviewed for the degree of thermal artefact and the margin status. Follow-up data after a 6-month-period were correlated with the margin status.

Results : Thermal artefact was present in all cases (mild 51.5%, moderate 36.2%, and severe 12.3%). In only one case, histologic diagnosis of the lesion was not possible due to severe thermal artefact. Nine cases (5.5%) had non-evaluable margins due to either thermal artefact (7 cases) or improper orientation of fragmented tissue (2 cases). Of 90 cases with subsequent surgical specimens, residual diseases were present in 4 of 21 (19.0%) with negative LEEP margins, in 31 of 64 (48.4%) with positive margins, and in 4 of 5 (80.0%) with non-evaluable margins.

Conclusions : Pathologic evaluation of the specimens from LEEP was limited in only a minority of cases. Thermal artefact was not a critical disadvantage of LEEP. The positive or negative margin status was correlated with the risk of residual disease.

Published in : J Med Assoc Thai 2001;84:507-514

CYTOLOGY OF SMALL-CELL CARCINOMA OF THE UTERINE CERVIX IN SEROUS EFFUSION: A REPORT ON TWO CASES

Surapan Khunamornpong *, Sumalee Siriaunkgul *, Prapaporn Suprasert **

* Department of Pathology

** Department of Obstetrics and Gynecology, Faculty of Medicine, Chiang Mai University

Cytologic features of 2 cases of small-cell carcinoma of the uterine cervix in the body fluid are described. Case 1 was a 34-yr-old woman with a stage IIA cervical tumor. Pleural effusion developed 6 months after initial diagnosis. Case 2 was a 38-yr-old woman with a stage IB tumor. Ascites was detected 11 mo after hysterectomy. Histologically, both cervical tumors were indistinguishable from small-cell carcinoma of oat-cell type in the lungs or other sites. Cytologically, the tumor cells in the pleural effusion of case 1 had characteristic features of small-cell carcinoma, including nuclear molding. However, almost all tumor cells in the ascites of case 2 showed a single-cell pattern mimicking malignant lymphoma. Mitotic figures and karyorrhetic bodies were occasionally seen. Nuclear molding was rarely identified. Small-cell carcinoma should be included in the differential diagnosis of malignant effusions containing lymphoma-like cells.

Published in: Diagn. Cytopathol. 2001; 24: 253-255.

PHASE II TRIAL OF DOCETAXEL AND CARBOPLATIN IN CISPLATIN-RECURRENT ADVANCED OVARIAN CANCER :A PRELIMINARY REPORT

Prapaporn Suprasert, Jatupol Srisomboon, Chailert Phongnarisorn, Chalong Cheewakriangkrai, Sithicha Siriaree, Sumitra Thongprasert*.

Division of Gynecologic Oncology, Department of OB & GYN,
* Division of Medical Oncology, Department of Medicine, Faculty of Medicine,
Chiang Mai University

Background : Patients with recurrent ovarian cancer have a poor prognosis with low response rates and short survival after salvage chemotherapy. In view of encouraging antitumor activity of docetaxel in the preclinical and phase I studies and the synergistic antitumor effect of this agent when combined with carboplatin, we accordingly, conducted a phase II trial of this regimen in patients with cisplatin-recurrent advanced ovarian cancer.

Objective : To evaluate the response rate and toxicity of docetaxel and carboplatin in patients with cisplatin-recurrent advanced epithelial ovarian cancer (EOC).

Methods : Twenty patients with cisplatin – recurrent EOC were recruited between October 1999- November 2001. Eligibility criteria included measurable > 2 cm. in diameter EOC. Age > 18 years old, an ECOG performance status of 0-2, adequate hepatic, renal and bone marrow function, and written informed consent. Docetaxel 75 mg/m² as 1 hour IV infusion and carboplatin AUC 5 were given on day 1 every 3 weeks for a maximum 6 cycles. All patients were evaluated for response, progression free survival (PFS) and WHO grading toxicity.

Results: Sixteen patients had recurrence more than 6 months (range 8-24 months) after cisplatin. Three patients had disease progression during receiving of cisplatin and 1 had recurrence within 6 months after receiving cisplatin. Four patients were withdrawn from the study because of moderated hypersensitivity (1 case), refusal of chemotherapy due to febrile neutropenia (2), and loss to follow up after 4 cycles (1 case). Fifteen patients were assessable for response. Eight patients had progression of disease. The overall response rate was 7 of 15 (46.7%) with 1 CR and 6 PR, all were the platinum sensitive patients. The median PFS was 9 months (95% CI = 6-12). Of the 88 cycles, grade 3 or 4 leukopenia and neutropenia occurred in 63.7% and 84.1%, respectively. Febrile neutropenia was found in 29.5%. Other toxicity was minimal.

Conclusion: The combination of docetaxel and carboplatin appear to have moderate efficacy and manageable toxicity in patient with cisplatin-recurrent epithelial ovarian cancer.

(Abstract#897). 38th Annual meeting : American Society of Clinical Oncology (ASCO) Conference. Orlando, Florida,USA, 19 May,2002.

THE ROLE OF EXTRAPERITONEAL PELVIC LYMPHADENECTOMY IN MANAGEMENT OF EARLY-STAGE CERVICAL CANCER: CHIANG MAI EXPERIENCE.

Jatupol Srisomboon*, Chailert Phongnarisorn*, Prapaporn Suprasert*, Kittipat Charoenkwan*, Chalong Cheewakriangkrai*, Sithicha Siriaree*, Patchaneeporn Porapakkham*, Aree Pantasart*, Sumalee Siriaunkgul**, Surapan Khunamornpong**

* Division of Gynecologic Oncology, Department of OB & GYN

** Division of Gynecologic Pathology, Department of Pathology, Faculty of Medicine, Chiang Mai University.

Lymph node metastases is the most important prognostic factor of early-stage cervical cancer after treatment by radical hysterectomy and pelvic lymphadenectomy (RHPL). For patients who have negative lymph nodes, the 5-year survival rate is 85-95% compared with 40-60% of patients with positive lymph nodes.

Surgical evaluation of pelvic lymph nodes prior to radical surgery could be performed via either transperitoneal or extraperitoneal approach. However, in patients receiving radiation therapy (RT) following transperitoneal pelvic lymphadenectomy (TPL), bowel complications (30%) are significantly higher than those (2.5%) receiving RT after extraperitoneal pelvic lymphadenectomy (EPL). In cases with grossly enlarged lymph nodes, radical hysterectomy should be abandoned to minimize serious morbidity occurring after combined treatment with radiation.

Our experience with EPL in management of early-stage cervical cancer are as follows
1. Stage IA2 cervical cancer patient undergoing inadvertent simple hysterectomy. The incidence of lymph node metastases is 5-7% and parametrial involvement is only 0.5%. We report 1 patient undergoing only EPL without morbidity.

2. Stage IB cervical cancer patients undergoing inadvertent simple hysterectomy. The incidence of lymph node metastases is 15-25% and parametrial involvement is 15-30%. We report 2 patients undergoing EPL followed by radical parametrectomy and upper vaginectomy without serious complications.

3. Stage IB – IIA cervical cancer. 30 patients treated with EPL prior to radical hysterectomy with uneventful consequence are presented. Radical hysterectomy was abandoned and treatment plan was switched to radiation therapy in 20% of cases by using this approach.

In conclusion, EPL could be safely performed before radical surgery or radiation for early-stage cervical cancer.

Presented at : 7th Annual Meeting of The Thai Gynecologic Oncology Group. Montien Hotel Pattaya, Thailand, August 10 – 12, 2002.

INVASIVE CERVICAL CANCER IN HUMAN IMMUNODEFICIENCY VIRUS INFECTED WOMEN IN CHIANGMAI, THAILAND

Vicharn Lorvidhaya, Pathomphorn Siraprasari, Prapaporn Suprasert,* Pimkhuan Kamnerdsupaphon

Division of Therapeutic Radiology and Oncology, Department of Radiology,
* Department of Obstetrics and Gynecology, Faculty of Medicine,
Chiang Mai University

Purpose : To evaluate treatment results of radiation, concurrent chemoradiation and the characteristics of invasive cervical carcinoma in human immunodeficiency virus (HIV) infected women.

Methods and materials : This is a descriptive analysis of 70 HIV – seropositive women with histologically confirmed invasive carcinoma of cervix. The documentation of HIV infection was confirmed with 2 times ELISA tests. All patients were clinically staged by gynecologic oncologist and radiation oncologist according to the FIGO staging system. Between January 1995 and December 2000 only 54 patients completed the planned treatments, which were radiotherapy and chemoradiotherapy. The mean follow – up time was 10.3 months, range 0.4 – 55.8. The Kaplan – Meier method was used for survival and disease – free survival analysis.

Results : The median age of the 70 patients was 38 years, range 22 – 61 years. The histology of sixty – three patients were squamous cell carcinomas, 4 were adenocarcinomas, 2 were adenosquamous cell carcinomas, and 1 was small cell carcinoma. Twelve (17%) patients were stage IB, 3(4.3%) were stage IIA, 27(38.6%) were stage IIB, 25(35.7%) were stage IIB, and 3(4.3%) were stage IVB. The 2 – year disease – free survival rate was 33.17%. The median time to progression was 9.5 months. We also founded that the patients with pretreatment lymphocyte counts more than 1,500 cells per cubic millimeters represented the better overall survival (p – value 0.042).

Conclusion : HIV infected patients represented a unique subset of cervical carcinoma and required the unique management. Radiation therapy and chemotherapy may be inadequate for these patients. Invasive cervical carcinoma in HIV infected women were diagnosed in younger age group, were in more advanced stage and had more rapid progression than immunocompetent patients. The degree of immunosuppression (pretreatment lymphocyte counts less than 1,500 cells per cubic millimeters) had significant effect on survival.

Presented at : 26th Annual Scientific Meeting on Mahidol's Day of The Faculty of Medicine, Chiang Mai University, Chiang Mai, September 24, 2002 (abstract page 36)

METASTATIC OR RECURRENT CERVICAL CANCER TREATED BY CISPLATIN PLUS 5-FU

Vicharn Lorvidhaya, Pimkhuan Kamnerdsupaphon, Prapaporn Suprasert.*

Division of Therapeutic Radiology and Oncology, Department of Radiology,
*Department of OB & GYN, Faculty of Medicine, Chiang Mai University

Objective : To assess the efficacy of chemotherapy with cisplatin and 5FU combination in patients with metastatic or recurrent cervical cancer.

Methods : Between October, 1999 and September 2001, 11 patients with metastatic/recurrent disease were enrolled to receive cisplatin : 80 – 100 mg/m² on day 1 in combination with 5FU : 800-1,000 mg/m² per day on day 1 through 4. Chemotherapy was administered on 3 week – schedule. Median age was 49 years (range 35 – 59); median ECOG performance status was 1.

Results : All eleven patients were evaluable for efficacy and safety. A total of 55 cycles were given with a median number of 6 cycles (range 4 – 6). Grade 3 anemia was recorded in 1 patient (9.1%). No patients experienced grade 3 – 4 neutropenia and infection. The main nonhematologic toxicity was nausea/vomiting but grade 3 was observed in only 1 patient (9.1%). Three patients achieved a complete response (27.3%) and 5 patients a partial response (45.5%); the overall response rate was 72.8%. The median progression – free survival was 8 months (7 – 16) with the median follow – up time of 10 months (4 – 17).

Conclusion : Our data confirm the efficacy of cisplatin+ 5FU regimen in cervical cancer. On this basis, phase III study is required to compare this regimen with others for this tumor.

Presented at : 26th Annual Scientific Meeting on Mahidol's Day of The Faculty of Medicine, Chiang Mai University, Chiang Mai, September 24, 2002 (abstract page 37)

RADIOCHEMOTHERAPY FOR LOCALLY ADVANCED SQUAMOUS VULVA CARCINOMA

Vicharn Lorvidhaya, Pimkuan Kamnerdsupaphon, Prapaporn Suprasert *

Division of Therapeutic Radiology and Oncology, Department of Radiology,
* Department of OB & GYN, Faculty of Medicine, Chiang Mai University**Objective :** To evaluate a regimen of radiochemotherapy as an alternative treatment for those patients with locally advanced squamous cell carcinoma of the vulva.**Methods :** Between July 1999 and February 2002, there were 11 patients with the diagnosis of squamous cell carcinoma of the vulva. Nine patients were stage IVA at presentation and 2 patients presented with stage III. All patients had biopsy prior to the treatment, and were treated with radiochemotherapy. Treatment included two cycles of chemotherapy with mitomycin - c (12 mg/m² on day 1) and fluorouracil (5FU 800 - 1,000 mg/m² on days 1 through 4) in addition to radiotherapy. Total radiation doses to the vulva and groins ranged from 45 to 70 Gy (median = 60), with pelvic doses of 45 to 60 Gy (median = 50).**Results :** There was a 100% overall response rate with complete responses in 7 patients (63.6%). Moist desquamation was the most important acute complication and required treatment interruptions in 2 patients. Among patients with complete clinical response, there has been 1 recurrence with the median follow-up of 8 months (2-21), and the time to progression was 5 months for this patient.**Conclusion :** This experience suggests that initial treatment with radiation and chemotherapy may offer some patients with locally advanced squamous cancer of the vulva an alternative to exenterative surgery and may hold curative potential for some patients with surgically unresectable of medically inoperable tumor.**Presented at :** 26th Annual Scientific Meeting on Mahidol's Day of The Faculty of Medicine, Chiang Mai University, Chiang Mai, September 24, 2002 (abstract page 38)**EVALUATION OF SAFETY AND EFFICACY OF TTS-FENTANYL IN ADULT PATIENTS WITH GYNECOLOGICAL CANCER - RELATED PAIN**Kanyarat Katanyoo, Vicharn Lorvidhaya, Jatupol Srisomboon, *
Prapaporn Suprasert*Division of Therapeutic Radiology and Oncology,
* Department of Obstetrics and Gynecology, Faculty of Medicine,
Chiang Mai University**Background :** Fentanyl, a synthetic strong opioid, has been recently incorporated into the Transdermal Therapeutic System [TTS]. The advent of a rate-limiting membrane provides constant release of the opioid. This trial was designed to establish the analgesic safety and efficacy of TTS - fentanyl in the treatment of chronic gynecologic cancer - related pain.**Objective :** To assess the analgesic safety and efficacy of TTS - fentanyl in the treatment of chronic gynecologic cancer - related pain.**Patients and Methods :** Thirty eligible patients were recruited into the study. Mean age was 47 years old. This open study comprise two phase. Phase 1 : an oral morphine stabilization phase, eligible patient who took other opioids and/or analgesic drugs were entered into the stabilization phase and should be converted to oral morphine according to conversion chart. The patients was then titrated to a stable oral morphine dose. Phase 2 : an open TTS - fentanyl treatment phase, the daily dose of oral morphine was switched to TTS - fentanyl according to conversion chart. The efficacy parameters are pain score assessments by visual analogue scale [VAS] and global assessments. The safety was evaluated by monitoring the patient's clinical conditions and adverse events.**Results :** TTS - fentanyl was generally well tolerated. Only 1 patient was terminated due to TTS - fentanyl treatment use because of adverse events. The most frequency was nausea or vomiting [46%] and constipation [33%]. The median pain VAS during TTS - fentanyl treatment decreased from 8 to 3 and global assessments at the end of treatment was better than the start of treatment.**Conclusion :** The results suggest that TTS - fentanyl is safe and effective in managing chronic gynecologic cancer - related pain.**Presented at :** 26th Annual Scientific Meeting on Mahidol's Day of The Faculty of Medicine, Chiang Mai University, Chiang Mai, September 24, 2002 (abstract page 40)**TECHNIQUE AND APPLICATION OF EXTRAPERITONEAL PELVIC LYMPHADENECTOMY IN CERVICAL CANCER**Jatupol Srisomboon, Chailert Phongnarisorn, Prapaporn Suprasert,
Kittipat Charoenkwan, Sitthicha Siriaree, Chalong Cheewakriangkrai,
Patchaneeporn Porapakkhham.Division of Gynecologic Oncology, Department of OB & GYN, Faculty of Medicine,
Chiang Mai University

Transperitoneal pelvic lymphadenectomy has been reported to be hazardous for cervical cancer patients who subsequently require radiation therapy when compared with those who undergo extraperitoneal pelvic lymphadenectomy (EPL). In addition to minimizing adhesion formation from peritoneal injury, extraperitoneal approach provides exposure of the retroperitoneal space, pelvic vessels and ureter while having an intact peritoneum to aid in retraction.

EPL can be performed via either midline or Maylard incision. The rectus muscles are bluntly dissected from the underlying peritoneum to enter the extraperitoneal space of the pelvis. The inferior epigastric vessels are clamped, cut and ligated. The Retzius space and paravesical space are developed by blunt dissection. The peritoneum is freed from iliac vessels and Cooper ligament by using blunt dissection with either the hand or a spongostick. Dissection is continued until obturator nerve is clearly visible deep in the pelvic side wall. The round ligament is then transected and ligated at its exit through the inguinal ring, this will aid in further exposure of the lateral space. The ureter is then easily identified on the peritoneum. A self-retaining retractor or 2 retractors are placed over the peritoneum and the patient is tilted toward the opposite side to aid further exposure of the pelvic vessels and obturator space. Pelvic node dissection could be carried out in a traditional fashion.

EPL could be applied in management of 1) stage IA2/IB cervical cancer patients undergoing inadvertent simple hysterectomy; 2) stage IA2 - IIA cervical cancer patients with planned Wertheim operation, if lymph node metastases is detected, radical hysterectomy should be abandoned; 3) cervical cancer patients prior to radiation therapy (surgical staging), if lymph node metastases is detected above the common iliac vessels, extended - field radiation should be instituted in addition to pelvic radiation.

Presented at 17th Annual Scientific Meeting of The Royal Thai College of OB & GYN. Lee Garden Plaza Hotel, Songkhla, Thailand, October 16-18, 2002.**SURGICAL EVALUATION OF PELVIC LYMPH NODES BY EXTRAPERITONEAL PELVIC LYMPHADENECTOMY BEFORE RADICAL HYSTERECTOMY FOR EARLY STAGE CERVICAL CANCER**Jatupol Srisomboon*, Patchaneeporn Porapakkhham*, Chailert Phongnarisorn *,
Prapaporn Suprasert *, Chalong Cheewakriangkrai *, Kittipat Charoenkwan *,
Sitthicha Siriaree*, Aree Pantusart*, Sumalee Siriaungkul**, Surapan Khunamornpong **.

*Division of Gynecologic Oncology, Department of OB & GYN

** Division of Gynecologic Pathology Department of Pathology, Chiang Mai University

Background : Lymph node metastases is the most important prognostic factor affecting survival of cervical cancer patients treated with radical hysterectomy. Complete removal of macroscopically positive nodes is of therapeutic advantage for patients compared to those who did not undergo resection of grossly positive nodes. Radiation - associated bowel complication could be dramatically reduced after node dissection was performed via extraperitoneal approach compared with transperitoneal approach.**Objective :** To determine the clinical benefit of extraperitoneal pelvic lymphadenectomy (EPL) prior to radical hysterectomy (RH) in early - stage cervical cancer.**Methods :** Between April 1, 2002, and June 30, 2002, 30 patients with stage IA - IIA cervical cancer underwent EPL before RH. Clinically enlarged nodes were completely resected and pathologically evaluated by frozen section. RH was abandoned in case of positive nodes. Perioperative data and early complications were recorded.**Results :** The patients were in stage IA with positive margins for invasive cancer (1), IB1 (20), IB2 (2), and IIA (7). The time from skin incision to complete EPL ranged from 45 mins, to 117 mins, with a mean of 65.5 mins. The total operative time ranged from 170 mins to 350 mins with a mean of 221 mins. The number of lymph nodes removed ranged from 7 - 55 with a mean of 28.5. The number of positive nodes ranged from 1 - 10 with a mean of 4.1. Macroscopically positive nodes were detected at EPL in 6 patients (20%) in whom RH was abandoned. Positive nodes were subsequently microscopically diagnosed in 3 patients (10%) who underwent EPL followed by RH. Three patients received adjuvant pelvic radiation for close surgical margins (2) and parametrial involvement (1). Of note is that 5 of 7 patients (71%) with stage IIA cervical cancer had positive pelvic nodes. Overall, 12 patients (40%) received postoperative radiation. No major complication was noted during EPL. Bladder injury occurred in 2 patients during RH**Conclusion :** EPL can be safely performed to clinically evaluate the lymph node status in cervical cancer patients prior to radical hysterectomy. Treatment plan was switched to radiation therapy in 20% of patients by using this approach.**Presented at :** 17th Annual Scientific Meeting of The Royal Thai College of OB & GYN. Lee Garden Plaza Hotel, Songkhla, Thailand, October 6 - 18, 2002.

PREVIOUS HYSTERECTOMY IN PATIENTS WITH OVARIAN CANCER : A 14 – YEAR REPORT FROM CHIANG MAI UNIVERSITY

Kittipat Charoenkwan, Jatupol Srisomboon, Prapaporn Suprasert, Chailert Phongnarisorn, Sithicha Siriaree, Chalong Cheewakriangkrai, Aree Pantusart.

Division of Gynecologic Oncology, Department of OB & GYN, Faculty of Medicine, Chiang Mai University

Objective : To determine the impact of prophylactic oophorectomy as a preventive measure for ovarian cancer in Thai women.

Methods : Medical records of 1,058 women with ovarian cancer evaluated and treated at Chiang Mai University Hospital between January 1988 and December 2001 were reviewed in an attempt to identify those who had hysterectomy performed with conservation of one or both ovaries at any point prior to diagnosis of ovarian cancer. Pertinent clinical and pathologic data of the women in this subset were further reviewed in detail.

Results : Of 1,058 women, 19 women (1.8%) had hysterectomy performed prior to diagnosis of ovarian cancer with 3 women (0.28%) had hysterectomy done at the age of 45 or above. The mean interval between the hysterectomy and the diagnosis of ovarian cancer was 8.6 years (range 7 months – 25 years). Considering the rate of prior hysterectomy at age 45 or beyond of 0.28% and the annual national ovarian cancer incidence of 1,252 cases, an estimated 3 – 4 cases of ovarian cancer could be prevented annually if prophylactic bilateral oophorectomy at the time of hysterectomy were routinely offered in this region.

Conclusions : The role and impact of incidental oophorectomy at the time of hysterectomy in general population as a preventive measure for subsequent ovarian cancer appeared to be significantly less prominent in this region of the world compared to Western or industrialized countries.

Presented at : 17th Annual Scientific Meeting of The Royal Thai College of OB & GYN. Lee Garden Plaza Hotel, Songkhla, Thailand, October 16 – 18, 2002.

WELL-DIFFERENTIATED VILLOGLANDULAR ADENOCARCINOMA OF THE UTERINE CERVIX : CYTOMORPHOLOGIC OBSERVATION OF FIVE CASES

Surapan Khunamornpong, Sumalee Siriaungkul, Prapaporn Suprasert*

Department of Pathology, Faculty of Medicine, Chiang Mai University

* Division of Gynecologic Oncology, Department of OB & GYN, Chiang Mai University

We retrospectively examined the cytologic findings of well-differentiated villoglandular adenocarcinoma (VGA) treated in our hospital. Cervical smears of four cases and a touch preparation of another case of VGA formed the materials for the study. The cytologic features were correlated with the histomorphology of VGA. Architecturally, long slender papillae and cohesive branching epithelial sheets with smooth borders and a lack of feathery edge were observed. Crowding and overlapping of nuclei were noted. The nuclei were uniform, small, and round to oval-shaped, with evenly distributed granular chromatin. Nucleoli were absent or inconspicuous. Mitoses were occasionally seen in all but one case. As the features of VGA are distinctive, the diagnosis could be possible on cytological grounds. Examination of cervical smears would be helpful for an early diagnosis of VGA or to suggest the coexistence of other neoplastic components.

Published in: Diagn Cytopathol 2002; 26: 10-14.

THE NECESSITY OF ROUTINE HEMOGLOBIN CHECK-UP IN CERVICAL CANCER PATIENTS RECEIVING RADIATION THERAPY.

Patchaneeporn Porapakham*, Bandit Chumworathayi*, Jarawan Tantipalakorn*, Prapaporn Suprasert*, Vicharn Lorvidhaya**, Jatupol Srisomboon*, Kittipat Charoenkwan*, Chailert Phongnarisorn*, Sithicha Siriaree*, Chalong Cheewakriangkrai*

* Division of Gynecologic Oncology, Department of OB & GYN

** Division of Therapeutic Radiology and Oncology, Chiang Mai University

Objective : To evaluate the value of weekly hemoglobin check up in cervical cancer patients receiving radiation therapy.

Design : Descriptive analysis.

Setting : Division of Gynecologic Oncology, Department of Obstetrics & Gynecology and Division of Therapeutic Radiology and Oncology, Department of Radiology, Faculty of Medicine, Chiang Mai University.

Methods : The medical records of cervical cancer patients admitted in gynecologic oncology ward between October 2001 and May 2002 were reviewed to evaluate the clinical characteristics and weekly hemoglobin level.

Results : There were 58 cervical cancer patients receiving radiation therapy in the study period. The median age was 56.9 years (range 25 – 81). Distribution by FIGO staging was stage I 3 (5.2%), II 21 (36.2%), III 29 (50%), and IV 1 (1.7%). Twenty – four (43.1%) patients had baseline hemoglobin less than 10 gm/dl. Among these patients, grade 2 – 4 anemia was found in 6 – 15 patients per week with a mean of 9.2 (38.5%) in week 1 – 4. Between week 5 – 8, grade 2 – 4 anemia was found in 4 – 10 patients per week with a mean of 6 (25%). Among 34 patients with based line hemoglobin \geq 10 gm/dl, grade 2 – 4 anemia was found in 4 – 6 patients per week with a mean of 5.2 (15.4%) in the first 4 weeks. After week 4, grade 2 – 4 anemia was found in only 2 – 3 patients per week with a mean of 2.3 (6.8%).

Conclusion : For cervical cancer patients with baseline hemoglobin less than 10 gm/dl, weekly hemoglobin level should be checked during radiation therapy. Among those with baseline hemoglobin more than 10 gm/dl, weekly hemoglobin check – up may not be necessary.

Presented at : 17th Annual Scientific Meeting of The Royal Thai College of OB & GYN. Lee Garden Plaza Hotel, Songkhla, Thailand, October 16 – 18, 2002.

SCRAPE CYTOLOGY OF THE OVARIES : POTENTIAL ROLE IN INTRAOPERATIVE CONSULTATION OF OVARIAN LESIONS.

Khunamornpong S, Siriaungkul S.

Department of Pathology, Faculty of Medicine, Chiang Mai University

Intraoperative diagnosis of ovarian lesions can be achieved by gross examination with the help of frozen sections and/or cytologic examination. Various cytologic techniques, including imprint, fine-needle aspiration, and scrape, may be used. In this study, we evaluated the application of scrape cytology in the diagnosis of ovarian lesions occurring during a 16-mo period at one institution. The cytologic diagnosis was primarily based on findings in air-dried, Diff-Quik-stained smears in correlation with clinical and intraoperative findings. In total, 131 histologically proven ovarian lesions, including 13 nonneoplastic lesions, 47 benign tumors, 17 epithelial tumors of low malignant potential (LMP), and 54 malignant tumors (35 primary, 1 leukemia, and 18 metastases), were cytologically examined. The accuracy of scrape cytology was 95 % in the benign, 47 % in the LMP, and 98 % in the malignant group. In the LMP group, the false-negative rate was 12 %, while the deferred rate and false-positive rate were 24 % and 18 %, respectively. There was no misdiagnosis between the benign and malignant categories. The histologic subtypes were correctly predicted in 78 % of cases. There were limitations of scrape cytology in the diagnosis of LMP and mucinous tumors, which require histologic architectural evaluation and adequate histologic sampling. Scrape cytology is a simple, rapid, and inexpensive adjunctive technique that should be used in intraoperative consultation for ovarian lesions.

Published in: Diagn Cytopathol 2003; 28(5): 250-7.

PREVALENCE AND PREDICTING FACTORS FOR PELVIC LYMPH NODE METASTASIS IN STAGE IB1 CERVICAL CARCINOMA.

Piyarat Udomwan,* Kittipat Charoenkwan,* Sumalee Siriakungul,** Jatupol Srisomboon,* Surapan Khunamornpong,** Prapaporn Suprasert*

* Department of Obstetrics & Gynecology

** Department of Pathology Faculty of Medicine, Chiang Mai University

Objective : To determine the prevalence of pelvic nodal metastasis and predicting factors in women with stage IB1 carcinoma of the cervix who underwent radical hysterectomy and pelvic lymphadenectomy.

Design : Cross sectional analytic study

Setting : Department of OB & GYN, Faculty of Medicine, Chiang Mai University

Subjects : Patients with stage IB1 carcinoma of the cervix who underwent radical hysterectomy and pelvic lymphadenectomy at Chiang Mai University Hospital between January 2000 and December 2002.

Results : Of 251 evaluable patients, radical hysterectomy was performed in 232 patients (92 %) and was abandoned in 19 patients (8 %). The prevalence of lymph node metastasis was 24 %. No clinical factors were significantly associated with pelvic lymph node metastasis. Histologic grade, depth of invasion, lymphovascular space invasion, parametrial invasion, and uterine invasion were significantly correlated with pelvic lymph node metastasis. From logistic regression analysis, the independent predicting factors for pelvic nodal metastasis were lymph vascular space invasion ($p < 0.0001$) and parametrial invasion ($p = 0.02$).

Conclusion : The prevalence of pelvic lymph node metastasis in stage IB1 carcinoma of the cervix is 24 %. The strongest predicting factors of node metastasis are lymphovascular space invasion and parametrial invasion.

Presented at : The 18th Annual Scientific Meeting of The Royal Thai College of OB & GYN, The Royal Golden Jubilee Building, Bangkok Thailand, 15-17 October, 2003 (abstract page 182)

Published in : Thai J Obstet Gynaecol 2003 ; 15 : 161 – 167.

THE FREQUENCY AND OUTCOME OF ABANDONED RADICAL HYSTERECTOMY IN CHIANG MAI UNIVERSITY HOSPITAL.

Prapaporn Suprasert,* Jatupol Srisomboon,* Kittipat Charoenkwan,* Sumalee Siriakungul,** Surapan Khunamornpong,** Sithicha Siriaree,* Chailert Phongnarisorn,* Thitima Siriwaranya,* Chalong Cheewakriangkrai.,* Vicharn Lorvidhaya**

* Division of Gynecologic Oncology, Department of OB & GYN

** Department of Pathology,

*** Division of Therapeutic Radiology and Oncology, Department of Radiology, Faculty of Medicine, Chiang Mai University

Objective : To evaluate the frequency and outcome of the abandoned radical hysterectomy (RH) in patients with cervical cancer.

Design : Retrospective study

Setting : Division of Gynecologic Oncology, Department of OB & GYN, Faculty of Medicine, Chiang Mai University.

Material & Methods : Between December 2000 and January 2003, the medical records of stage IA2-IIA cervical cancer patients planned to undergo radical hysterectomy and pelvic lymphadenectomy (RHPL) were reviewed. The clinical characteristics and disease free survival (DFS) in patients whose plan for radical hysterectomy was abandoned due to positive nodes were compared with those who had positive nodes after RHPL.

Results : There were 242 patients undergoing RHPL in the study period. Of these patients, 23 cases (9 %) were found at the operation to have grossly positive pelvic nodes. The radical hysterectomy was abandoned and only pelvic and paraaortic lymph node dissection were carried out. Six patients (26.1 %) received neoadjuvant chemotherapy. Twenty-two (95.6 %) patients subsequently received concurrent chemoradiation. One patient received radiation alone. Extended field radiation of paraaortic area were added in 4 (17.4 %) cases who had positive paraaortic nodes. When compared with 62 patients who had positive nodes after RHPL, there were statistically significant difference in terms of histology, number of positive nodes and tumor size. The adenocarcinoma was more frequent in group of abandoned-RH (34 % vs 11 %, $P = 0.01$). The mean number of positive nodes (5 vs 1, $P = 0.01$) and tumor size (3.1 vs 2.1 cm, $P = 0.02$) in the abandoned RH group were also higher than those of the RHPL group (21 % vs 11 %, $P = 0.01$). The 2-year DFS was nearly statistically significant lower in the abandoned-RH group when compared with the RHPL group (68 % vs 93 %, $P = 0.06$).

Conclusion : Larger tumor size, number of positive nodes, adenocarcinoma histology, and complication are significantly more common in the abandoned-RH patients. The DFS of these patients is lower than those whose positive nodes are detected after RHPL.

Presented at : The 18th Annual Scientific Meeting of The Royal Thai College of OB & GYN, The Royal Golden Jubilee Building, Bangkok Thailand, 15-17 October, 2003 (abstract page 247)

OUTCOME OF HIGH-RISK EARLY STAGE CERVICAL CANCER TREATED WITH RADICAL HYSTERECTOMY AND PELVIC LYMPHADENECTOMY.

Thitima Siriwaranya,* Prapaporn Suprasert,* Sumalee Siriakungul,** Surapan Khunamornpong,** Jatupol Srisomboon,* Kittipat Charoenkwan,* Sithicha Siriaree,* Chailert Phongnarisorn,* Chalong Cheewakriangkrai,* Vicharn Lorvidhaya.***

* Department of Obstetrics & Gynecology,

** Department of Pathology,

*** Division of Therapeutic Radiology and Oncology, Department of Radiology, Faculty of Medicine, Chiang Mai University

Objective : To evaluate the survival and complications of high-risk early stage cervical cancer treated with radical hysterectomy and pelvic lymphadenectomy.

Design : Retrospective descriptive study.

Setting : Division of Gynecologic Oncology Department of Obstetrics and Gynecology, Faculty of Medicine, Chiang Mai University.

Subject : 157 patients with FIGO stage IA2 to IIA cervical cancer treated between January 1998 to January 2003, undergoing radical hysterectomy with pelvic lymphadenectomy and had at least one high risk pathological factors.

Intervention : Pelvic lymphadenectomy was initially performed followed by radical hysterectomy. Patients with high risk pathological factors were treated with adjuvant radiation with or without concurrent chemotherapy.

Result : With median follow up of 26 months, the projected 5-year progression-free survival and overall survival were 80.0 % and 92.8 % respectively. Treatment-related complication occurred in 15 patients (9.5 %), all of these patients received adjuvant radiation. The incidences of lymph node metastasis and parametrial involvement were 18.8 % and 11.2 % respectively. Tumor recurrence occurred in 12 patients (7.6 %) in which 8 of these has lymph node metastasis, the remaining had parametrial invasion.

Conclusion : Early stage cervical cancer patients with high-risk pathological factors had relatively high survival rate with low complication in this setting.

Presented at : The 18th Annual Scientific Meeting of The Royal Thai College of OB & GYN, The Royal Golden Jubilee Building, Bangkok Thailand, 15-17 October, 2003 (abstract page 214)

Published in : Thai J Obstet Gynaecol 2003 ; 15 : 93 – 9.

CONCURRENT MITOMYCIN C, 5-FLUOROURACIL, AND RADIOTHERAPY IN THE TREATMENT OF LOCALLY ADVANCED CARCINOMA OF THE CERVIX : A RANDOMIZED TRIAL

Vicharn Lorvidhaya,* Imjai Chitapanarux,* Supatra Sangruchi,** Prasert Lertsanguansinchai,*** Yongyut Kongthanarat,+ Saipin Tangkaratt,++ Ekachai Visetsiri+++

* Faculty of Medicine, Chiang Mai University

** Siriraj Hospital, Mahidol University,

*** Chulalongkorn Hospital, Chulalongkorn University

+ Rajvithi Hospital,

++ National Cancer Institute,

+++ Bhumibol Adulyadej Hospital

Purpose : This is a prospective, Phase III multicenter randomized trial to assess the effectiveness of concurrent intravenous mitomycin C, oral 5-fluorouracil (5-FU), and radiotherapy (RT) in locally advanced carcinoma of the cervix.

Methods and Materials : Between January 1988 and November 1994, 926 patients with locally advanced carcinoma of the cervix, FIGO Stage IIB-IVA, were entered into this study. The patients were randomized into four arms, as follows : Arm 1 : conventional RT; Arm 2 : conventional RT and adjuvant chemotherapy; Arm 3 : conventional RT plus concurrent chemotherapy; Arm 4 : conventional RT plus concurrent chemotherapy and adjuvant chemotherapy. Concurrent chemotherapy consisting of intravenous mitomycin C at 10 mg/m² was given on Days 1 and 29, and oral 5-FU at 300 mg/day was administered on Days 1-14 and 29-42 during RT. Adjuvant chemotherapy of 5-FU orally at 200 mg/day was given for three courses of 4 weeks, with a 2-week rest every 6 weeks. Six centers participated in the trial.

Results : The median follow-up time was 89 months. Acute side effects were generally higher in concurrent arms, but most of the patients tolerated the treatment well. Bone marrow toxicity was also higher in concurrent arms. The 5-year actuarial disease-free survival (DFS) was 48.5%, 54.1%, 64.5% and 59.7% for arms 1,2,3, and 4, respectively. The pattern of failure revealed a significant increase in locoregional recurrence in the nonconcurrent chemotherapy arm. The local recurrence was 25.5%, 20.6%, 14.3%, and 17.6% for arm 1,2,3, and 4, respectively. The metastatic rates were not significantly different in all four arms. At the time of analysis, there were no increases in late side effects, especially in gastrointestinal and genitourinary systems.

Conclusions : Concurrent chemotherapy, mitomycin C, and 5-FU together with conventional RT showed an improved DFS rate when compared with conventional RT alone in patients with locally advanced carcinoma of the cervix.

Published in : Int J Radiation Oncology Biol Phys 2003; 55(5) : 1226-32.

COMPARISON OF ORAL VERSUS INTRAVENOUS RAMOSETRON IN PREVENTION OF ACUTE CISPLATIN – INDUCED EMESIS : A RANDOMIZED CONTROLLED TRIAL.

Charuwan Tantipalakov*, Jatupol Srisomboon, * Hunsu Thienthong**, Aree Pantusart, * Prapaporn Suprasert, * Chanram Saeresongkhun, ** Saisunee Eamtang, ** Amphika Sutthichat, ** Huthaya Pautad**

* Department of OB&GYN, Faculty of Medicine, Chiang Mai University, ** Division of Nursing, Maharaj Nakorn Chiang Mai Hospital, Chiang Mai

Objective : To compare the antiemetic efficacy of a single oral versus intravenous (IV) ramosetron, a new class of selective 5 – HT₃ receptor antagonists, in gynecologic cancer patients receiving high – dose cisplatin.

Method : Between February 2003 and July 2003, 109 patients with gynecologic cancer scheduled to receive single agent cisplatin chemotherapy at a dose of 75 mg/m² were randomized to receive either 0.2 mg oral (51 cases) or 0.3 mg IV (58 cases) ramosetron 1 h and 30 min respectively before chemotherapy. Patients were evaluated for 24 h after chemotherapy. The number of nausea and vomiting including adverse events were recorded every 6 h.

Results : 51 and 58 patients received oral and IV ramosetron respectively. Both groups were similar regarding age, performance status, body mass index and diagnosis of gynecologic cancer. 95 percent of cases were cervical cancer. Antiemetic effect was significantly higher in the oral group when compared with the IV group during the first 6 hours and during the period of 18 to 24 hours after administration of cisplatin chemotherapy. Overall in 24 h, patients receiving oral ramosetron experienced no emesis slightly higher than that of the IV group (55% and 36% respectively, p = 0.05). Adverse events were mild and transient and were not significantly different in both groups, except tiredness which was more frequent in the IV group.

Conclusion : Oral ramosetron at a dosage of 0.2 mg is as effective as 0.3 mg of intravenous ramosetron in prevention of acute emesis in patients receiving 75 mg/m² of cisplatin chemotherapy.

Published in : J Med Assoc Thai 2004 ; 87 : 119 – 125.

Presented at : The 56th Annual Scientific Meeting of Japan Society of Obstetrics and Gynecology, Tokyo, Japan, April 10 – 14, 2004.

ROLE OF PROPHYLACTIC OOPHORECTOMY AT THE TIME OF HYSTERECTOMY IN OVARIAN CANCER PREVENTION IN THAILAND.

Kittipat Charoenkwan, Jatupol Srisomboon, Prapaporn Suprasert, Chailert Phongnarisorn, Sitthicha Siriaree, Chalong Cheewakriangkrai.

Division of Gynecologic Oncology, Department of Obstetrics and Gynecology, Faculty of Medicine, Chiang Mai University

Aim : To determine the impact of prophylactic oophorectomy on ovarian cancer prevention in Thai women by estimating the magnitude of reduction in ovarian cancer incidence in Thailand if the procedure was routinely offered.

Methods : A database of 752 women with epithelial ovarian cancer treated at Chiang Mai University Hospital between January 1990 and December 2001 was reviewed in an attempt to identify those who had hysterectomy performed for indications other than ovarian and corpus cancer with conservation of one or both ovaries prior to diagnosis of ovarian cancer. Pertinent clinical and pathologic data of the women in this subset were further reviewed in detail.

Results : Of 752 ovarian cancer patients, 13 (1.73 %) had hysterectomy performed prior to the diagnosis of ovarian cancer with only one woman having (0.13 %) had a hysterectomy at the age of 45 or above. The mean interval between the hysterectomy and the diagnosis of ovarian cancer was 9.9 years (range 2.25), considering the rate of prior hysterectomy at age 45 or beyond of 0.13 % and the annual national ovarian cancer incidence of 1252 cases, an estimated one to two cases of ovarian cancer could be prevented annually if prophylactic bilateral oophorectomy at the time of hysterectomy in a woman age 45 or beyond were routinely offered in Thailand.

Conclusion : Prophylactic oophorectomy at the time of hysterectomy in general Thai population as a preventive measure for subsequent ovarian cancer should not be recommended without complete knowledge of patients' socioeconomic background and propensity to comply with hormone replacement therapy regimens.

Published in : J Obstet Gynaecol Res 2004; 30(1) : 20-23.

PULMONARY METASTASES IN GESTATIONAL TROPHOBLASTIC TUMOR : 6 YEARS EXPERIENCE IN CHIANG MAI UNIVERSITY HOSPITAL

Prapaporn Suprasert¹, Juntima Eua – throngchit², Jatupol Srisomboon¹, Kittipat Charoenkwan¹, Sitthicha Siriaree¹, Chailert Phongnarisorn¹

¹Division of Gynecologic Oncology, Department of OB&GYN

²Division of Diagnostic Radiology, Department of Radiology, Faculty of Medicine, Chiang Mai University

Objective : To evaluate the patterns and clinical outcome of pulmonary metastases in patients with gestational trophoblastic tumor (GTT).

Methods : The medical records and chest films of GTT patients receiving treatment at Chiang Mai University Hospital between January 1998 and June 2003 were reviewed.

Results : There were 85 GTT patients in the study period. Of these patients 32 cases (37.6%) had pulmonary metastases. The mean age was 30 years (21 ~ 55 years). The previous pregnancy were abortion, term pregnancy and molar in 8 (25%), 10 (31.3%) and 14 (43.8%) respectively. The median interval between the previous pregnancy and development of GTT was 24 months (1 ~ 288 months). The median initial B – hCG level was 106,537 IU/L (range 140 – 2,557,500 IU/L). The metastases confined to the lungs alone in 10 cases (31.2%), and the remaining 22 patients had metastases elsewhere also, i.e. pelvic organs, brain, bladder, kidney, vagina, liver and spleen. Nine (28.1%) cases presented with lung symptoms. The most common pulmonary metastases pattern was multiple lung infiltration (16), followed by large and solitary nodule (11), pleural effusion (3), tiny infiltration (1) and atelectasis (1). Twenty – seven patients (84.4%) received multiple chemotherapy and 13 (40.6%) required more than one regimens. The mean number of chemotherapy was 7 cycles (range 3 ~ 23). Three patients received whole brain radiation for brain metastasis. Hysterectomy was performed in 12 (37.5%). Thoracotomy and craniotomy were each done in 2 patients. One case received chest radiation. Among 10 patients with only lung metastases, 8 (80%) had complete response with no recurrence, the remaining 2 patients were lost to follow – up. Among 22 patients with multiple organ metastases, 16 (72%) had complete response, 2 died from disease, 4 were lost to follow – up.

Conclusion : The most common pattern of pulmonary metastases was multiple lung infiltration followed by solitary nodule and pleural effusion. Patients with only lung metastases had better prognosis than those with lung and other organ metastases.

Presented at : The 56th Annual Congress of the Japan Society of Obstetrics and Gynecology, LeMeridien Grand Pacific Hotel, Tokyo, Japan, April 11 – 13, 2004. * Received President Award of the JSOG

THE INCIDENCE OF CERVICAL INTRAEPITHELIAL NEOPLASIA BY CONTRACEPTIVE METHOD IN A COHORT OF THAI WOMEN.

Gupta SB, Srisomboon J, Liaw K, Wootipoom V, Go V, Yuenyao P, Vipupinyo C, Celentano D

Objective : To determine the incidence of cervical intraepithelial neoplasia (CIN) in women using different contraceptive methods.

Methods : This on – going prospective study enrolled 1255 Thai women aged 18 – 35 to examine the natural history of HPV infection and cervical neoplasia. Women are screened for cervical abnormalities every 6 months using the ThinPrep liquid – based cytology technique. Women with abnormal ThinPrep results undergo colposcopy, and subsequently, cervical biopsy if a lesion is visible under the colposcope.

Results : At the baseline visit, 1110/1255 (88%) women enrolled in the study had normal cytology results, including 324 (29%) oral contraceptive (OC) users, 426 (38%) injectable depot medroxyprogesterone acetate (DMPA) users, and 360 (32%) users of non – hormonal methods. The overall incidence of CIN 1 and CIN 2/3 in the cohort at the 6 – month visit was 4.07 and 1.11 per 100 person – years (PYs), respectively. The incidence of CIN 1 at the 6 – month visit was 0.74, 2.22, and 1.11 per 100 PYs in OC, DMPA, and non – hormonal method users, respectively. The incidence of CIN 2/3 was 0.92, 0, 0.18 per 100 PYs in OC, DMPA, and non – hormonal method users, respectively.

Conclusions : Compared to estimated incidence rates of cervical neoplasia in the literature, the incidence of CIN 1 and CIN 2/3 at the 6 – month follow – up visit in this cohort of Thai women is high. Although the incidence of CIN 1 was disproportionately higher in the DMPA group compared to users of other contraceptive methods, the incidence of CIN 2/3 was not higher in this group. More follow – up data are needed to examine whether there are true differences in the incidence of cervical neoplasia based on the contraceptive use.

Presented at : 21st International Papillomavirus Conference, Mexico City, Mexico 2004.

NERVE – SPARING RADICAL HYSTERECTOMY. A NEW TREND IN SURGICAL TREATMENT OF EARLY – STAGE CERVICAL CANCER TO REDUCE THE PELVIC AUTONOMIC NERVE INJURY : CHIANG MAI EXPERIENCE.

Kittipat Charoenkwan, Jatupol Srisomboon , Prapaporn Suprasert , Chailert Phongnarisorn, Sitticha Siriaree, Chalong Cheewakriangkrai, Charuwan Tantipalakorn , Chumnan Kietpeerakool.

Division of Gynecologic Oncology, Department of OB & GYN, Faculty of Medicine, Chiang Mai University.

Conventional radical hysterectomy for early cervical cancer has contributed a favorable outcome with 5 – year survival over 90% but is associated with typical long – term morbidity, i.e. bladder dysfunction, colorectal motility disorders and sexual dysfunction. Pelvic autonomic nerve injury during surgery is judged to be the main cause of these dysfunctions. Accordingly, more attempt has been made to minimize such morbidity after surgery, whilst the radicality of the parametrial resection is still maintained.

Nerve – sparing radical hysterectomy is the operation aimed to reduce pelvic organ dysfunctions by identifying and preserving the pelvic autonomic nerves during the parametrial resection. The operation can be summarized into 3 steps as follows :

Step 1 : Preservation of the hypogastric nerve and the proximal part of the inferior hypogastric plexus during the resection of the posterior parametrium or the sacrouterine and the rectouterine ligaments.

Step 2 : Preservation of the pelvic splanchnic nerves and the middle part of the inferior hypogastric plexus during the resection of the lateral parametrium or the cardinal ligaments.

Step 3 : Preservation of the distal part of the inferior hypogastric plexus during the resection of the anterior parametrium or the vesicouterine ligaments.

Ten patients with early – stage cervical cancer undergoing nerve – sparing radical hysterectomy in Chiang Mai University Hospital will be presented. The main outcome including feasibility, safety, and efficacy focusing on the postoperative voiding function, as well as the obstacles will be discussed.

Presented at : 9th Annual Scientific Meeting of The Thai Gynecologic Cancer Society, Golden Sand Resort Hotel, Petchburi, Thailand, August 12 – 14, 2004.

MALIGNANT OVARIAN GERM CELL TUMOR (MOGCT): EXPERIENCE IN CHIANG MAI UNIVERSITY HOSPITAL, THAILAND.

Prapaporn Suprasert, Jatupol Srisomboon, Chailert Phongnarisorn, Kittipat Charoenkwan, Sitticha Siriaree, Sumalee Siriaunkgul *, Surapan Khunamornpong *

Division of Gynecologic Oncology, Department of OB & GYN, * Department of Pathology, Faculty of Medicine, Chiang Mai University

Background & Aim : MOGCT are rare gynecologic tumors usually affecting young women. We conducted the retrospective study to evaluate the clinical feature and treatment outcome of this tumor.

Methods : The medical records of MOGCT patients receiving treatment at Chiang Mai University Hospital between 1998 and 2002 were reviewed.

Results : There were 461 ovarian cancer patients in the study period. Of these patients, 46 (9.9%) were MOGCT. The mean age was 23 years (7 – 39 years). The most common presenting symptom was abdominal mass (26) followed by abdominal enlargement (12), abdominal pain (7) and abnormal uterine bleeding (1). 22 patients (47.8%) underwent conservative surgery. The FIGO stage distribution was stage I (31), stage II (2), and stage III (11). Two patients presented with recurrence. 19 (41.3%) had immature teratoma, 11 (23.9%) had dysgerminoma, 11 (23.9%) had endodermal sinus tumor and 5 (10.9%) had mixed type. 30 (65.2%) were treated with bleomycin, etoposide and cisplatin (BEP). The overall response rate was 84.7% (complete response 80.4%, partial response 4.3%). 4 (8.7%) had progression of disease. 5 – year disease free survival was 89.6% Severe treatment related – complications were pulmonary fibrosis (2), septicemia (1), electrolyte imbalance (1) and bowel obstruction (2). Two patients died from toxicity of chemotherapy, 3 died from disease and 5 were lost to follow up.

Conclusion : The most common histology of MOGCT was immature teratoma. The outcome of this tumor was good although some had severe treatment – related complications.

Presented at : 10th Biennial International Gynecologic Cancer Society Meeting (IGCS), Edinburgh, Scotland October 3 – 7, 2004.

RUPTURED MATURE CYSTIC TERATOMAS MIMICKING ADVANCED STAGE OVARIAN CANCER : A REPORT OF 2 CASES STUDY.

Prapaporn Suprasert*, Surapan Khunamornpong**, Sumalee Siriaunkgul **, Chailert Phongnarisorn *, Sitticha Siriaree *

* Department of OB & GYN, Faculty of Medicine, Chiang Mai University
** Department of Pathology, Faculty of Medicine, Chiang Mai University

Spontaneous rupture of mature cystic teratoma is uncommon. Chronic rupture of teratomas may result in granulomatous peritonitis. In rare cases, the clinical presentation and the intraoperative findings may mimic those of advanced stage ovarian cancers. Two cases of ruptured mature cystic teratomas mimicking malignancy are reported in 50 – and 53 – year – old patients. Both presented with abdominal distension and weight loss. Elevation of serum CA – 125 (233 unit/ml) was detected in one case. Intraoperatively, nodular thickening of the omentum with marked peritoneal adhesion was observed in both cases. Ascites of 1500 ml was present in one case. Pathologic examination in both cases showed ovarian mature cystic teratomas with peritoneal granulomatous inflammations response to the released tumor content. Both patients were followed for 44 and 12 months and were free of symptoms without additional treatment other than surgery. Intraoperative pathologic consultation can help confirm the benign diagnosis and unnecessary major operation for malignancy can be avoided.

Published in : J Med Assoc Thai 2004 ; 87 (12) : 1522 – 1525

TREATMENT RESULTS OF METHOTREXATE AND FOLINIC ACID AS PRIMARY CHEMOTHERAPY FOR NONMETASTATIC GESTATIONAL TROPHOBLASTIC NEOPLASIA.

Jatupol Srisomboon *, Prapaporn Suprasert*, Chailert Phongnarisorn *, Kittipat Charoenkwan*, Sitticha Siriaree *, Chalong Cheewakriangkrai *, Charuwan Tantipalakorn *, Chumnan Kietpeerakool *

* Department of Obstetrics and Gynecology, Faculty of Medicine Chiang Mai University.

Objective : To evaluate the efficacy and toxicity of methotrexate and folinic acid (MTX–FA) chemotherapy in patients with nonmetastatic gestational trophoblastic neoplasia (NMGTN).

Patients and Methods : Between 1997 and 2003, a total of 67 patients with NMGTN received treatment at the Chiang Mai University Hospital. Of the 67 patients, 55 were initially treated with methotrexate 1.0 mg/kg intramuscularly (IM) on day 1, 3, 5 and 7 and folinic acid 0.1 mg/kg IM on day 2, 4, 6 and 8. Treatment courses were repeated every 14 days. Clinical characteristics and outcomes were analyzed.

Results : All 55 patients with NMGTN were cured. Of the 55 patients initially treated with MTX–FA, 49 (89%) achieved complete remission. Six (11%) patients developed methotrexate resistance, 3 were cured with actinomycin D, 1 were cured with 5–fluorouracil followed by etoposide, 2 required hysterectomy to attain remission. No serious toxicity was noted.

Conclusion : Methotrexate and folinic acid chemotherapy is highly effective and well – tolerated in treating patients with nonmetastatic gestational trophoblastic neoplasia.

Published in : J Med Assoc Thai 2005; 88: 886-90.

OUTCOMES OF ABANDONED RADICAL HYSTERECTOMY IN PATIENTS WITH STAGE IB – IIA CERVICAL CANCER FOUND TO HAVE POSITIVE NODES DURING THE OPERATION.

Prapaporn Suprasert*, Jatupol Srisomboon*, Kittipat Charoenkwan*, Sumalee Siriaunkgul**, Surapan Khunamornpong**, Sitthicha Siriaree*, Chailert Phongnarisorn*, Vicharn Lorvidhaya***

* Division of Gynecologic Oncology, Department of OB & GYN
 ** Department of Pathology,
 *** Division of Therapeutic Radiology and Oncology,
 Department of Radiology, Faculty of Medicine, Chiang Mai University

The objective of this study was to evaluate the outcomes of stage IB – IIA cervical cancer patients whose radical hysterectomy (RH) was abandoned for positive pelvic nodes detected during the operation compared with those found to have positive nodes after the operation. Among 242 patients with planned RH and pelvic lymphadenectomy (RHPL) for stage IB – IIA cervical cancer, 23 (9.5%) had grossly positive nodes. RH was abandoned, complete pelvic lymphadenectomy was performed. Of these 23 patients, 22 received adjuvant chemoradiation, the remaining 1 received adjuvant radiation. Four patients with positive paraortic nodes were additionally treated with extended field irradiation.

When compared with 35 patients whose positive nodes were detected after the operation, there were significant differences regarding number of positive nodes and number of patients receiving extended field irradiation. Complications in both groups were not significantly different, but the 2 – year disease – free survival was significantly lower in the abandoned RH group compared with that of the RHPL group (58.5% vs 93.5%, $P = 0.01$). In conclusion, the survival of stage IB – IIA cervical cancer patients whose RH was abandoned for grossly positive pelvic nodes was significantly worse than that of patients whose node metastasis was identified after the operation. This is because the abandoned RH group had worse prognostic factors.

Published in : Int J Gynecol Cancer 2005; 15: 498-502.

RADICAL HYSTERECTOMY FOR STAGE IIB CERVICAL CANCER : A REVIEW.

Prapaporn Suprasert, Jatupol Srisomboon, Takahiro Kasamatsu.*

Division of Gynecologic Oncology, Department of OB & GYN,
 Faculty of Medicine, Chiang Mai University.

* Department of Gynecology, National Cancer Center Hospital, Tokyo 104, Japan.

Patients with stage IIB cervical cancer in some countries in Europe and Asia especially in Japan are usually treated with radical hysterectomy and pelvic lymphadenectomy. Extrauterine disease, i.e. nodal metastases, parametrial invasion and intraperitoneal spread can be readily identified. We present the literature review of radical hysterectomy in stage IIB cervical cancer by searching data since 1980 from MEDLINE and found that the parametrial involvement of patients in this stage was only 21 – 55%, the incidence of pelvic node metastases was about 35 – 45%, and 5 – year survival was between 55 – 77%. Lymph node metastases and the number of positive nodes were significant prognostic factors of patients in this stage.

Published in: Int J Gynecol Cancer 2005; 15: 995-1001.

METASTATIC TUMORS TO THE OVARIES: A STUDY OF 170 CASES IN NORTHERN THAILAND.

Surapan Khunamornpong¹, Prapaporn Suprasert², Wittanee Na Chiangmai,³ Sumalee Siriaunkgul¹

¹Department of Pathology, Faculty of Medicine, Chiang Mai University,
²Department of Obstetrics and Gynecology, Faculty of Medicine,
 Chiang Mai University

³Department of Radiology, Faculty of Medicine, Chiang Mai University

The cases of malignant ovarian tumors treated at Chiang Mai University hospital between 1992 and 2003 were histologically reviewed. The medical records, the radiologic findings, and the follow-up outcome in the cases suspicious or diagnostic of metastases were reviewed to confirm the diagnosis and to determine the primary sites. Metastatic tumors accounted for 30 % of malignant ovarian tumors. A total of 170 cases of metastatic tumors included 117 cases with non-gynecologic origins and 53 cases with gynecologic origins. Non-gynecologic metastatic tumors were from large intestine (31 %), stomach (14 %), intrahepatic bile duct (10 %), breast (9 %), extrahepatic bile duct/gallbladder (7 %), appendix (5 %), hematologic tumors (3 %), others (4 %), and unknown primary site (16 %). Metastatic gynecologic tumors were from cervix (53 %), corpus (34 %), fallopian tube (11 %), and gestational trophoblastic disease (2 %). The proportion of metastatic tumors to malignant ovarian tumors in northern Thailand was comparable to those of the Western or Japanese studies. However, the distribution of the primary sites was different and was correlated with the cancer incidence in Thai women. The majority of mucin-producing adenocarcinomas involving the ovaries were metastatic tumors.

Published in : Int J Gynecol Cancer 2006 ;16 Suppl 1:132-8.

CLEAR CELL ADENOCARCINOMA OF THE FEMALE GENITAL TRACT : PRESENCE OF HYALINE STROMA AND TIGROID BACKGROUND IN VARIOUS TYPES OF CYTOLOGIC SPECIMENS

Surapan Khunamornpong¹, Paul S. Thorne², Prapaporn Suprasert³, Sumalee Siriaunkgul¹

¹Department of Pathology, Faculty of Medicine, Chiang Mai University,

²Department of Laboratory Medicine, Hospital for Sick Children and University of Toronto, Toronto, Canada

³Department of Obstetrics and Gynecology, Faculty of Medicine,
 Chiang Mai University

Hyaline basement membrane-like stromal material and tigroid background are distinctive cytologic features in Diff-Quik- or Giemsa-stained smears of clear cell adenocarcinoma (CCA) of the female genital tract. However, it is uncertain how often these findings are present in different types of cytologic specimens, and which type of preparation is optimal for making this diagnosis. We therefore reviewed the cytologic features of CCA in 3 types of specimens including scrape cytology (15 specimens), fine needle aspiration (FNA) (7 specimens), and peritoneal cytology (15 specimens), with emphasis on the findings in Diff-Quik-stained smears. The cell morphology in scrape cytology and FNA was comparable, whereas in peritoneal cytology, the cytoplasm was better preserved. Most tumor cells had fragile cytoplasm containing variable amounts of fine vacuoles, and round nuclei with distinct or prominent nucleoli. Hyaline stroma was present in 93% of scrape cytology specimens, 71% of FNA specimens, and 80% of peritoneal cytology specimens. Tigroid background was observed in 47% of scrape cytology specimens, 43% of FNA specimens, but in none of the peritoneal cytology specimens. Formation of a tigroid background may be prevented by the abundant fluid content in this type of cytologic specimen. Hyaline stroma and tigroid background were uncommonly seen in scrape smears from other types of primary ovarian tumors, mainly juvenile granulosa cell tumor and yolk sac tumor. However, the additional presence of papillary structures allows CCA to be readily distinguished from these other tumours. Scrape cytology would be the best method to choose for the intraoperative cytologic diagnosis of CCA.

Published in: Diagn Cytopathol 2005; 32: 336-40.

YOLK SAC TUMOR OF THE VULVA: A CASE REPORT WITH LONG-TERM DISEASE-FREE SURVIVAL

Surapan Khunamornpong^a, Sumalee Siriaungkul^a, Prapaporn Suprasert^b,
Imjai Chitapanarux^c

^a Department of Pathology, Faculty of Medicine, Chiang Mai University,

^b Department of Obstetrics and Gynecology, Faculty of Medicine,
Chiang Mai University

^c Division of Therapeutic Radiology and Oncology, Faculty of Medicine,
Chiang Mai University

Background. Yolk sac tumor (YST) of the vulva is extremely rare. Seven cases of vulvar YST have been reported to the literature. Due to the rarity of tumors, the appropriate choice of treatment may remain unclear.

Case. A 30-year-old woman presented with a 3.5-cm right labial mass. Excisional biopsy showed YST with predominant solid pattern. Three weeks after excision, right inguinal lymph node biopsy revealed metastatic tumor. The serum alpha-fetoprotein (AFP) was not elevated. Cisplatin-based chemotherapy was administered, followed by pelvic and groin irradiation. The patient was free of disease 90 months after the diagnosis.

Conclusion. Local excision of tumor with adjuvant cisplatin-based chemotherapy can be justified for vulvar YST. Inguinal lymphadenectomy is recommended because metastasis may occur early. Adjuvant radiation therapy may help to control the disease. Tumor size of 5 cm or less may be a favorable prognostic factor. Serum AFP level may not be a sensitive marker for follow-up of vulvar YST.

Published in: *Gynecol Oncol* 2005; 97: 238-242.

SURVIVAL AND PROGNOSTIC FACTORS FOR PATIENTS WITH EARLY-STAGE CERVICAL CANCER TREATED WITH RADICAL SURGERY : STAGE IB1 VS. IB2.

Jatupol Srisomboon*, Kittipat Charoenkwan*, Sumalee Siriangkul**,
Surapan Khunamornpong**, Prapaporn Suprasert*, Chailert
Phongnarisorn*, Sitthichai Siriaree*, Chalong Cheewakriangkrai*.

* Division of Gynecologic Oncology, Department of OB & GYN

** Department of Pathology, Faculty of Medicine, Chiang Mai University

Objective: To compare the clinical outcome of stage IB1 vs. stage IB2 cervical cancer patients treated with radical hysterectomy and pelvic lymphadenectomy (RHPL).

Methods: Patients with stage IB cervical cancer undergoing RHPL at Chiang Mai University Hospital between January 1998 and December 2002 were evaluated for survival and recurrence. Clinicopathological variables were analyzed to identify the prognostic factors affecting survival and recurrence.

Results: During the study period, RHPL was performed on 432 stage IB1 and 14 stage IB2 patients. The incidence of positive pelvic nodes were 17.8% and 28.6% for stage IB1 and IB2 respectively ($p = 0.29$). Incidence of parametrial involvement were similar between the stages IB1 and IB2 (10.9% and 14.3%) respectively, $p = 0.66$. Patients with stage IB2 received adjuvant radiation more frequently than IB1 patients (64.3% vs 35.9%, $p = 0.03$). Overall, 19/442 patients (4.3%) developed recurrence, 4% for stage IB1, and 14.3% for stage IB2 ($p = 0.12$). With a median follow-up of 37.5 months, the 3-year disease-free survivals were 96% and 80% for stage IB1 and stage IB2, respectively ($p = 0.03$).

Conclusion: Patients with stage IB1 cervical cancer appear to have significantly better outcome than those with stage IB2.

Presented at: The 57th Annual Congress of The Japan Society of Obstetrics and Gynecology, Kyoto International Conference Hall, Kyoto, Japan, April 2-5, 2005.

ADVERSE AFFECTS OF PACLITAXEL AND CARBOPLATIN COMBINATION CHEMOTHERAPY IN EPITHELIAL GYNECOLOGIC CANCER.

Chumnan Kietpeerakool, Prapaporn Suprasert, Jatupol Srisomboon

Department of Obstetrics and Gynecology, Faculty of Medicine,
Chiang Mai University,

Objective: To evaluate the adverse affects of paclitaxel and carboplatin combination chemotherapy.

Subjects: Patients with epithelial cancer of the ovary, fallopian tube and peritoneum treated with paclitaxel and carboplatin combination chemotherapy at Chiang Mai University Hospital between August 2003 and August 2004.

Results: Of 224 evaluable cycles in 63 patients treated with paclitaxel (175 mg/m²) and carboplatin (AUC 5), grade 3 and 4 neutropenia occurred in 37.1% or 41.3% of patients. 4.8% of patients experienced febrile neutropenia. Grade 3 and 4 leukopenia occurred in 8.6% of courses and 12.6% of patients. Grade 3 anemia occurred in 5.2% of courses and 9.5% of patients. Grade 3 thrombocytopenia occurred in 2.8% of courses and 9.6% of patients. The nonhematologic adverse affects were rare, however some adverse events may be potentially life threatening.

Conclusion: Adverse affects of paclitaxel and carboplatin combination chemotherapy are acceptable and manageable in the majority of patients.

Published in: *J Med Assoc Thai* 2005; 88: 301-6.

PRIMARY CARCINOMA OF THE FALLOPIAN TUBE : A CLINICOPATHOLOGIC ANALYSIS OF 27 PATIENTS.

Chumnan Kietpeerakool, Prapaporn Suprasert, Jatupol Srisomboon,
Aree Pantusart.

Department of OB & GYN, Faculty of Medicine, Chiang Mai University

Objectives: To analyze the clinicopathologic features of women with primary fallopian tube carcinoma.

Design: Descriptive cross sectional study.

Material and Method: Twenty-eight women diagnosed with primary fallopian tube carcinoma treated at Chiang Mai University Hospital between January 1997 and December 2004.

Results: During the study period, the primary fallopian tube carcinoma accounted for 0.48% of all gynecologic malignancies. Of the 28 patients, one was excluded for unavailable medical records. Mean age at diagnosis was 53 years (range, 38-76 years). Seventeen (63.0%) were menopausal women. The most common clinical presentation was pelvic mass (55%), followed by abnormal vaginal bleeding (18.5%). Hydrops tubae profluens was present in three (11.1%) women. The rare presenting symptoms including pelvic peritonitis and abnormal glandular cells on cervicovaginal smear were noted in one (3.7%) woman of each category. In all women, primary fallopian tube carcinoma could not be diagnosed preoperatively. During the operation, an abnormal tubal lesion was suspected in only eleven (40.7%) women. Histology were serous adenocarcinoma (70.4%), endometrioid adenocarcinoma (22.2%), undifferentiated adenocarcinoma (3.7%) and carcinosarcoma (3.7%). As opposed to epithelial ovarian cancer, the majority of women in the present study were in the early stage of the disease.

Conclusion: Primary fallopian tube carcinoma is a rare gynecologic malignancy that has various and nonspecific presentations. Definite diagnosis is usually made postoperatively. This malignancy should be considered in differential diagnosis of peri- and postmenopausal women who present with complex adnexal mass, unexplained uterine bleeding, abnormal glandular cells on cervicovaginal smear and complicated pelvic inflammatory disease.

Published in: *J Med Assoc Thai* 2005; 88 (10): 1338-43.

CLINICOPATHOLOGIC PREDICTORS OF INCOMPLETE EXCISION AFTER LOOP ELECTROSURGICAL EXCISION PROCEDURE FOR CERVICAL NEOPLASIA

Chumnan Kietpeerakool, Jatupol Srisomboon, Kannaporn Ratchusiri.

Department of Obstetrics and Gynecology, Faculty of Medicine, Chiang Mai University, Chiang Mai

The aim of this study was to identify the factors affecting incomplete excision after loop electrosurgical excision procedure (LEEP) for evaluation and treatment of cervical neoplasia. Patients with abnormal cervical cytology who underwent colposcopy and LEEP at Chiang Mai University Hospital between October 2004 and July 2005, were retrospectively evaluated. During the study period, 201 patients were eligible for analysis. All cone margin involvement was observed in 44% of the patients (95% CI, 37.3-51.4). Multivariate analysis revealed that invasive cancer on cytology (adjusted odds ratio [aOR] =3.05, 95% confidence interval [CI] =1.03 to 9.00; P=0.02), invasive cancer on LEEP histopathology (aOR=9.73, 95% CI =3.95 to 23.9; P<0.001), and cone length of less than 10 mm (aOR =1.95, 95% CI =1.04 to 3.66; P =0.03) were significant predictors for any cone margin involvement. For endocervical margin involvement, postmenopausal status and cone length of less than 10 mm were significant predictors of incomplete excision. As opposed to endocervical margin involvement, postmenopausal status was significantly associated with a decreased risk of ectocervical margin involvement. Invasive cancer on histopathology was a significant predictor of both ecto- and endocervical margin involvement. In Conclusions, Invasive cancer either on cytology or LEEP specimens and cone length of less than 10 mm are significant predictors of incomplete excision.

Keywords: Loop electrosurgical excision procedure, cone margin, cervical neoplasia, multivariate analysis.

Published in : Asian Pacific Journal of Cancer Prevention 2005; 6(4): 481-4

TOTAL LAPAROSCOPIC HYSTERECTOMY IN CERVICAL CANCER STAGE IA1 OR PERSISTENT HIGH GRADE SQUAMOUS INTRAEPITHELIAL LESIONS AFTER PREVIOUS DIAGNOSTIC CONIZATION.

Phongnarisorn C, Charoenkwan K, Srisomboon J, Uttavichai C.

Department of OB & GYN, Faculty of Medicine, Chiang Mai University

Objective : To describe preliminary experience on total laparoscopic hysterectomy (TLH) in the patients with cervical cancer stage IA1 or persistent high grade squamous intraepithelial lesion (HSIL), who underwent diagnostic cervical conization.

Design : A university teaching hospital.

Method and subjects : From June 2003 to September 2004, twenty patients with history of at least one previous diagnostic conization were recruited for TLH in our hospital. Of them, 16 has stage IA1 cervical cancer and the remaining 4 had persistent HSIL.

Results : TLH was successfully performed in all cases without conversion. The mean age of the patients was 46.5 ± 8.8 years (range 35-63). The mean body mass index was 23.8 ± 4.0 (range 18.5 - 33.7). The mean operation time was 271.4 ± 62.8 minutes (range 153-450). The mean uterine weight was 81.8 ± 42.1 grams. (range 20-190). The average time of postoperative hospital stay was 3 days. The mean pre- and post-operative hemoglobin were 12.4 g% and 10.3 g% respectively. Major complications occurred in two patients, cystotomy requiring laparoscopic repair in one case and ureteral injury requiring subsequent laparotomy in the other. Notably, these complications occurred in the first 4 cases of our series, at the beginning of our learning curve. However, both were healthy and had no significant morbidity at one-year of follow-up.

Conclusion : Based on this series, TLH for cervical carcinoma stage IA1 or persistent HSIL in case of previous conization may probably be safe and can be performed as an alternative to conventional laparotomy.

Presented at : The 14th Annual Congress of the International Society for Gynecologic Endoscopy, Hilton London Metropole, London, United Kingdom, April 3-6, 2005.

CONCURRENT CISPLATIN-BASED CHEMORADIATION AND ADJUVANT HYSTERECTOMY FOR BULKY STAGE IB-IIA CERVICAL CANCER.

Chalong Cheewakriangkrai**, Jatupol Srisomboon**, Prapaporn Suprasert**, Chailert Phongnarisorn**, Imjai Chitapanarux*, Sitthicha Siriaree**, Kittipat Charoenkwan**.

* Division of Therapeutic Radiology and Oncology, Department of Radiology;

** Division of Gynecologic Oncology, Department of Obstetrics and Gynecology, Faculty of Medicine, Chiang Mai University

Objective : To evaluate the outcomes and adverse effects of concurrent cisplatin-based chemoradiation and adjuvant hysterectomy for bulky stage IB-IIA cervical cancer.

Methods : All eligible, thirty-four patients with bulky stage IB-IIA cervical cancer were assigned to receive weekly cisplatin 40 mg/m² for 6 cycles concurrently with radiation followed by extrafascial hysterectomy 6 weeks after completion of radiation.

Results : Estimated 5-year progression-free and disease-free survival rates of 80% were observed after a median follow-up of 42 months. The overall recurrent rate was 18%. Grade 3 neutropenia and anemia were noted in only 5.9% and 2.9%, respectively. All acute toxicities were transient and manageable. There were no treatment-related deaths or late toxicities.

Conclusion : For appropriately selected patients with bulky stage IB-IIA cervical cancer, concurrent cisplatin-based chemoradiation followed by adjuvant hysterectomy offers an effective treatment option with acceptable toxicity.

Presented at : The 57th Annual Congress of The Japan Society of Obstetrics and Gynecology, Kyoto International Conference Hall, Kyoto, Japan, April 2-4, 2005.

OUTCOME OF INTERMEDIATE RISK FACTORS IN EARLY STAGE CERVICAL CANCER.

Prapaporn Suprasert, Jatupol Srisomboon, Surapan Khunamornpong*, Sitthicha Siriaree, Kittipat Charoenkwan, Chalong Cheewakriangkrai.

Division of Gynecologic Oncology, Department of OB&GYN
*Division of Gynecologic Pathology, Department of Pathology
Faculty of Medicine, Chiang Mai University

Objective : To evaluate the clinical outcome of cervical cancer patients who had intermediate risk factor(s) after treatment with radical hysterectomy.

Design : Retrospective study

Setting : Division of Gynecologic Oncology, Department of OB & GYN, Faculty of Medicine, Chiang Mai University.

Material & Methods : From January 1999 to January 2004, the medical records of cervical cancer patients who had intermediate risk factors defined as positive lymphovascular space invasion (LVSI) and/or deep stromal invasion (DSI) after treatment with radical hysterectomy and pelvic lymphadenectomy (RHPL) were reviewed. Patients who had LVSI more than 10 spaces per HPF with DSI less than 3mm. from cervical serosa received adjuvant chemoradiation. While patients who had LVSI > 10 spaces per HPF alone were treated with adjuvant chemotherapy (cisplatin).

Results : There were 154 patients with intermediate risk factors in the study period. The mean age was 43 years (28-67 years). Most of them were in stage IB1 (120 or 77.9%). The most common histology was squamous cell carcinoma (107 or 69.5%). The distribution of patients by histologic grade (G) was 40 in G1, 73 in G2 and 20 in G3, 21 were unclassified grade. The mean number of nodes removed was 28 nodes (11-70). 109 (70.8%) patients had positive LVSI. Of these patients, 38 had LVSI > 10 spaces. 92 patients (60%) had DSI. The treatment consisted of RHPL alone (90 or 58.4%), RHPL + chemoradiation (23 or 14.9%), RHPL + radiation (12 or 7.8%) and RHPL + chemotherapy (29 or 18.8%). The 5-year disease free survival (DFS) was 91.8%. Eleven patients (7.1%) developed recurrence 8-37 months after completion of treatment. Three patients had lung metastases, 2 after RHPL alone, the remaining 1 after adjuvant chemoradiation. One patient developed liver metastasis after adjuvant chemotherapy. Seven patients developed pelvic recurrence, 3 after RHPL alone, 3 after adjuvant chemotherapy and 1 after adjuvant radiation. Neither LVSI nor DSI was the significant prognostic factor when analyzed by univariate and multivariate analysis.

Conclusion : The outcome of early-stage node-negative cervical cancer who had LVSI and/or DSI after RHPL was quite good. Neither of them was the significant prognostic factors.

Presented at : The 57th Annual Congress of The Japan Society of Obstetrics and Gynecology, Kyoto International Conference Hall, Kyoto, Japan, April 2-4, 2005.

A RARE FEMALE GENITAL TRACT TUMOR : BENIGN GRANULAR CELL TUMOR OF VULVA : CASE REPORT AND REVIEW OF THE LITERATURE

Chalong Cheewakriangkrai^a, Sameer Sharma^a, George Deeb^b, Shashikant Lele^{a,*}.

^a Department of Gynecologic Oncology, Roswell Park Cancer Institute, Elm and Carlton Streets, Buffalo, NY 14263, USA.

^b Department of Pathology, Roswell Park Cancer Institute Buffalo, NY 14263, USA.

Background : The granular cell tumor (GCT) of vulva is a rare female genital tract tumor.

Case : A 59-year-old female was incidentally noted to have a 1-cm sized lump in her left labium mass that subsequently increased in size to approximately 4 cm. The mass was totally excised under general anesthesia. Her postoperative follow-up was uneventful. Histologic examination of the tumor showed sheets and clusters of infiltrating tumor cells with morphologic features consistent with granular cell tumor.

Conclusion : Although benign and slow growing, it has a tendency for recurrence and can cause morbidity and mortality when presenting with multicentric or multiple organ involvement due to the lack of effective systemic therapy. Therefore, clinicians and pathologists should be aware of its clinical and histopathologic features.

Published in : *Gynecol Oncol* 2005; 97 : 656-8.

EXTENT OF LYMPHOVASCULAR SPACE INVASION AND RISK OF PELVIC LYMPH NODE METASTASES IN STAGE IB1 CERVICAL CANCER.

Anchalee Chandacham*, Kittipat Charoenkwan*, Sumalee Siriaungkul**, Jatupol Srisomboon*, Prapaporn Suprasert*, Chailert Phongnarisorn*, Chalong Cheewakriangkrai*, Sithicha Siriaree*, Charuwan Tantipalakov*, Chumnan Kietpeerakool*.

* Department of OB & GYN, Faculty of Medicine, Chiang Mai University

** Department of Pathology, Faculty of Medicine, Chiang Mai University

Objective : To evaluate whether the extent of lymphovascular space invasion (LVSI) is a risk factor for pelvic lymph node metastases in stage IB1 cervical cancer.

Material and Method : The clinicopathological data of 397 patients with stage IB1 cervical cancer undergoing radical hysterectomy and pelvic lymphadenectomy (RHPL) at Chiang Mai University Hospital between January 1998 and December 2002 were analyzed. The histology, tumor grade, depth of stromal invasion, uterine corpus involvement, parametrial invasion and LVSI were analyzed for their association with pelvic node metastases. The extent of LVSI was classified as negative, minimal (< 10 LVSI / cervical specimen), and extensive (≥ 10 LVSI / cervical specimen).

Results : Of the 397 patients, 146 (36.8%) had tumors containing LVSI, 82 (20.7%) and 64 (16.1%) had minimal and extensive LVSI, respectively. Fifty nine (14.9%) patients had pelvic node metastases. Using multivariable analysis, LVSI ($p < 0.001$), depth of stromal invasion ($p < 0.001$), tumor grade ($p < 0.001$), and parametrial invasion ($p < 0.001$) were significant predictors of pelvic node metastases. The extent of LVSI either minimal or extensive degree significantly influenced pelvic node metastases.

Conclusion : The presence of LVSI especially extensive involvement was significantly associated with the risk of pelvic node metastases in stage IB1 cervical cancer.

Published in : *J Med Assoc Thai* 2005; 88 (Suppl 2): S31-6.

RADICAL SURGERY FOR T1 AND T2 SQUAMOUS CELL CARCINOMA OF THE VULVA THROUGH SEPARATE INCISIONS.

Apichart Khobjai, Jatupol Srisomboon, Kittipat Charoenkwan, Chailert Phongnarisorn, Prapaporn Suprasert, Chalong Cheewakriangkrai, Sithicha Siriaree, Charuwan Tantipalakov, Chumnan Kietpeerakool.

Division of Gynecologic Oncology, Department of Obstetrics and Gynecology, Faculty of Medicine, Chiang Mai University, Chiang Mai.

Objective : The aim of the study was to retrospectively evaluate treatment results in patients with T1 and T2 vulvar carcinoma.

Material and Method : The medical records of 46 patients with T1 and T2 SCC of the vulva undergoing radical excision of the tumor and groin node dissection at Chiang Mai University Hospital between January 1998 and December 2004 were reviewed. The tumor size, histologic grade, nodal status, lymph-vascular space invasion, lesion location, surgical marginal status, complications, recurrence and survival were analyzed.

Results : Mean age of the 46 patients (T1 = 15, T2 = 31) was 59 years with a range of 34-84 years. The incidence of lymph node metastases for T1 lesions was 13% compared to 35% for T2 lesions. Twenty nine patients (63%) experienced surgical complications, the most common one was lymphedema (16) while wound breakdown was noted in only 1 patient. With a median follow-up of 15 months, 14 patients (30%) developed recurrence, 3 (20%) and 11 (35%) in patients with T1 and T2 lesions respectively. The overall 5-year disease-free survival and 5-year survival were 37% and 40%, respectively. The 5-year survival of patients with T1 lesion was significantly higher than that of patients with T2 lesion (64% vs 31%, $P = 0.04$). Patients with negative nodes had significantly better survival than those with positive nodes (56% vs 18%, $P = 0.02$). In multivariable analysis, only the status of groin node remained an independent prognostic factor for survival.

Conclusion : Radical excision and groin node dissection through separate incision for T1 and T2 squamous cell carcinoma of the vulva in this study has a less favorable survival outcome compared with the literature.

Published in : *J Med Assoc Thai* 2005; 88 (Suppl 2) : S75-81.

RADIOLOGIC FEATURES AND TREATMENT OUTCOMES OF PULMONARY METASTASIS IN GESTATIONAL TROPHOBLASTIC NEOPLASIA

Prapaporn Suprasert*, Juntima Eua-throngchit**, Jatupol Srisomboon**, Kittipat Charoenkwan*, Sithicha Siriaree*, Chailert Phongnarisorn*.

* Department of OB & GYN, Faculty of Medicine, Chiang Mai University

** Department of Radiology, Faculty of Medicine, Chiang Mai University

Objective: To evaluate the radiologic patterns and treatment outcome of pulmonary metastasis in patients with gestational trophoblastic tumor (GTT).

Material and Method: The medical records and chest films of GTT patients treated at Chiang Mai University Hospital between January 1998 and June 2003 were reviewed.

Results: There were 85 GTT-patients in the study period. 32 cases (37.6%) had pulmonary metastasis diagnosed with chest X-rays. The most common radiologic pattern was well defined nodule. The radiologic features of patients who had lung metastases alone were not significantly different from those who had associated metastases in other organs. 27 patients (84.3%) received multiple chemotherapy and 6 required more than one regimen. The mean number of chemotherapy was 7 cycles (range 3-23). Adjuvant surgery consisted of hysterectomy (11), salpingo-oophorectomy (1), thoracotomy (2), and craniotomy (1). Four patients received whole brain irradiation for brain metastases. Among 10 patients with lung metastasis alone, 8 (80%) attained complete remission, the remaining 2 patients were lost to follow-up. Among 22 patients with associated multiple organ metastases, 16 (72.7%) had complete remission, 2 died from diseases, 4 were lost to follow-up.

Conclusion: The most common radiologic pattern of pulmonary metastasis in GTT patients was well-defined multiple lung nodules. The radiologic features of patients who had lung metastases alone were not significantly different from those who developed metastases in other organs.

Published in : *J Med Assoc Thai* 2005; 88(7): 875-80.

ACCURACY OF VISUAL INSPECTION WITH ACETIC ACID (VIA) TEST IN PREDICTING ASSOCIATED VAGINAL NEOPLASIA IN PATIENTS WITH EARLY STAGE CERVICAL CANCER UNDERGOING RADICAL HYSTERECTOMY

Prapaporn Suprasert, Jatupol Srisomboon, Kittipat Charoenkwan, Sithicha Siriaree, Chalong Cheewakriangkrai, Chumnan Kietpeerakool.

Division of Gynecologic Oncology, Department of OB & GYN, Faculty of Medicine, Chiang Mai University, Chiang Mai

Background: Associated vaginal neoplasia is found in about 8-10% of patients with early stage cervical cancer causing positive vaginal margins after radical hysterectomy and needs adjuvant brachytherapy. Vaginal neoplasia usually stains white after application of acetic acid. This study used the VIA test to identify concomitant vaginal neoplasia in the hope that it may reduce positive vaginal margins.

Objective: To evaluate the accuracy of VIA test in predicting associated vaginal neoplasia in patients with early stage cervical cancer undergoing radical hysterectomy.

Methods: Early stage cervical cancer patients with planned radical hysterectomy February 2004 and May 2005 in Chiang Mai University Hospital were tested with 5% acetic acid on the vagina and cervix during the preoperative pelvic examination. Patients with abandoned radical hysterectomy patients were excluded from the study. Positive VIA test was defined as acetowhite vaginal epithelium. The VIA test and pathology of the resected vagina were analyzed.

Result: There were 70 patients in the study period. 3 patients were excluded for abandoned radical hysterectomy. Of the 67 patients, 9 had positive VIA test but only 1 had concomitant vaginal neoplasia. The sensitivity, specificity, positive and negative predictive values of the VIA testing were 11.1%, 84.5, 10% and 85.9% respectively. 4 patients had positive vaginal margins for vaginal neoplasia.

Conclusion: VIA test is not sensitive in detecting associated vaginal neoplasia in patients with early-stage cervical cancer.

Presented: (Poster Number 94). The 14th International Meeting of the European Society of Gynaecological Oncology (ESGO). Hilton Istanbul Hotel, Istanbul, Turkey, September 28,2005

WEEKLY VERSUS THREE-WEEKLY CISPLATIN AS AN ADJUNCT TO RADIATION THERAPY IN HIGH-RISK STAGE I-IIA CERVICAL CANCER AFTER SURGERY: A RANDOMIZED COMPARISON OF TREATMENT COMPLIANCE.

Bandit Chumworathayi, Prapaporn Suprasert, Kittipat Charoenkwan, Jatupol Srisomboon, Chailert Phongnarisorn, Sitticha Siriaree, Chalong Cheewakriangkrai, Jaruwan Tantipalakorn, Chumnan Kiatpeerakool, Aree Pantusart.

Division of Gynecologic Oncology, Department of OB&GYN, Faculty of Medicine, Chiang Mai University, Chiang Mai.

Objective: To compare weekly and three-weekly cisplatin as an adjunct to radiation therapy in high-risk early-stage cervical cancer after surgery with regard to treatment compliance.

Material and Method: From June 1st, 2003 to February 29th 2004, we performed a randomized trial of radiotherapy in combination with two concurrent chemotherapy regimens, weekly or three-weekly cisplatin in patients with high-risk cervical cancer FIGO stage I-IIA after surgery. Women with primary invasive squamous-cell carcinoma adenocarcinoma or adenosquamous carcinoma of the cervix were enrolled. The patients also had to have an absolute neutrophil count of at least 1,500 cells per cubic millimeter, a platelet count of at least 75,000 cells per cubic millimeter, a creatinine clearance higher than 40 milliliter per minute, and adequate hepatic function. All patients received external-beam radiotherapy according to a strict protocol. Patients were randomly assigned to receive one of two chemotherapy regimens : 75 mg per square meter of cisplatin on days 1,22, 43 and 64 or every three weeks for 4 cycles (group 1) or 40 mg per square meter of cisplatin per week for six cycles (group 2).

Results: The analysis included 40 women. First group that received three-weekly cisplatin had a higher rate of incomplete and delayed treatments than the second group that received weekly cisplatin ($p < 0.001$ and $P = 0.0236$ respectively). The relative risks of delayed courses were 2.06 (95 percent confidence interval. 1.15 to 3.68) for group 1, as compared with group 2. The toxicity-related incomplete treatment rate and G-CSF doses used were significantly higher in group 1 than in group 2.

Conclusion: Concurrent chemoradiation with weekly cisplatin regimen has more complete treatment rate and less delayed courses than that with three-weekly cisplatin among women with high-risk cervical cancer after surgery.

Published in: J Med Assoc Thai 2005; 88 (11): 1483-1492.

CLINICAL OUTCOMES AND PROGNOSTIC FACTORS OF NODE-NEGATIVE CERVICAL CANCER PATIENTS WITH DEEP STROMAL INVASION OR LYMPHOVASCULAR SPACE INVOLVEMENT FOLLOWING RADICAL HYSTERECTOMY

Prapaporn Suprasert¹, Jatupol Srisomboon¹, Sumalee Siriaungkul², Surapan Khunamornpong², Chailert Phongnarisorn¹, Kittipat Charoenkwan¹, Sithichah Siriaree¹, Chalong Cheewakriangkrai¹, Chumnan Kietpeerakool¹

¹ Department of Obstetrics and Gynecology,

² Department of Pathology, Faculty of Medicine, Chiang Mai University, Chiang Mai

OBJECTIVE: To evaluate the clinical outcomes and prognostic factors of node-negative cervical cancer patients who had deep stromal invasion (DSI) and/or lymphovascular space invasion (LVSI) following radical hysterectomy and pelvic lymphadenectomy (RHPL).

MATERIAL AND METHOD: The medical records of 150 node-negative stage IA2-IIA cervical cancer patients who had DSI and/or LVSI after RHPL from 1999 to 2004 were reviewed.

RESULTS: Eighty-eight (58.4%) patients were treated with RHPL alone. Twenty-eight (18.7%), 23 (15.4%), eight (5.3%), and three (2%) patients received postoperative chemotherapy, chemoradiation, radiotherapy, and brachytherapy, respectively. Overall, 11 (7.3%) patients developed recurrence. The estimated 5-year disease-free survival of the patients was 90.9%. By multivariate analysis, two factors, age of less than 35 years old and a non squamous histology, were significantly independent prognostic. Eight (5.3%) patients experienced treatment-related complications.

CONCLUSION: Node-negative cervical cancer patients with DSI and/or LVSI had excellent clinical outcomes. Young age and non-squamous histology are significant independent prognostic factors.

Published in: J Med Assoc Thai 2006; 89(9): 1368-75.

THE RELATION OF OVARIAN CANCER AND ENDOMETRIOSIS

Prapaporn Suprasert¹, Surapan Khunamornpong²,

¹Department of Obstetrics and Gynecology, ²Department of Pathology, Faculty of Medicine, Chiang Mai University, Chiang Mai

This study was undertaken to evaluate the clinical features and survival outcomes of ovarian cancer patients who had associated pelvic endometriosis. The medical records of 1076 patients with ovarian cancer treated at Chiang Mai University Hospital between 1995 and 2005 were reviewed. Among of these patients, 37 (3.4%) had associated pelvic endometriosis. The mean age of the patients was 44 years (25-62 years). The most common presenting sign and symptom was abdominal mass (12), followed by abdominal pain (10), abdominal distension (7), abnormal uterine bleeding (2). Twenty-one (56.8%) patients were nulliparous and 14 (37.8%) were single. The stage distribution was stage I (24), stage II (4), stage III (4), and stage IV (1). Four patients had 2 primary carcinomas. The most common histology of the 37 patients was clear cell carcinoma (17) followed by endometrioid carcinoma (11). The estimated 5-year disease - free survival was 55.4%. In conclusion, most patients associated with endometriosis-associated ovarian carcinoma present with abdominal mass and abdominal pain. Clear cell CA is the most common histology in ovarian cancer patients who have associated endometriosis. Three fourths of the patients are in stage I and have favorable prognosis.

Published in: Asian Pac J Cancer Prev 2006; 7(4):

THE RISK OF RESIDUAL NEOPLASIA IN WOMEN WITH MICROINVASIVE SQUAMOUS CERVICAL CARCINOMA AND POSITIVE CONE MARGINS.

Chailert Phongnarisorn¹, Jatupol Srisomboon¹, Surapan Khunamornpong², Sumalee Siriaungkul², Prapaporn Suprasert¹, Kittipat Charoenkwan¹, Chalong Cheewakriangkrai¹, Sitthicha Siriaree¹, Tawiwat Pantasri¹.

¹ Departments of Obstetrics & Gynecology,

² Department of Pathology, Faculty of Medicine, Chiang Mai University,

The objective was to evaluate the prevalence and factors affecting residual disease in women with cervical microinvasive carcinoma (MIC) with positive cone margins for high-grade lesions and invasive carcinoma. We reviewed histopathology slides of 129 women with MIC who had high-grade lesions or invasive carcinoma at cone margins. These patients underwent hysterectomy following cone biopsy between January 1994 and June 2004. Of the 129 patients, 77 (59.7%) had residual disease in the hysterectomy specimens, in which 57 (44.2%) had residual high-grade lesions. Twenty patients (15.5%) had residual invasive carcinoma: 18 were microinvasive and 2 were invasive. Factors significantly affecting the risk of residual disease included positive postconization endocervical curettage ($P=0.001$), positive cone margins for invasive carcinoma ($P=0.003$), and depth of stromal invasion >1 mm ($P=0.014$). Cox proportional hazards analysis revealed positive cone margins for invasive carcinoma as significant predictor of residual invasive disease (hazard ratio, 3.22; 95% CI 1.21-8.60, $P=0.019$). In summary, patients with MIC and positive cone margins for high-grade lesions or invasive carcinoma are at high risk of residual neoplasia. Repeat cone biopsy should be performed to determine exactly the severity of lesion before planning treatment.

Published in: *Int J Gynecol Cancer* 2006; 16(2): 655-9.

HIGH-GRADE SQUAMOUS INTRAEPITHELIAL LESION WITH ENDOCERVICAL CONE MARGIN INVOLVEMENT AFTER CERVICAL LOOP ELECTROSURGICAL EXCISION: WHAT SHOULD A CLINICIAN DO ?

Sitthichai Siriaree¹, Jatupol Srisomboon¹, Chumnan Kietpeerakool¹, Sumalee Siriaungkul², Surapan Khunamornpong², Apaporn Natpratan³, Sumon Pratheapjarus³, Amornrat Futemwong³, Uraporn Chantarasenawong³

¹ Department of Obstetrics and Gynecology, ² Department of Pathology,

³ Obstetrics and Gynecologic Emergency Room Unit,

Faculty of Medicine, Chiang Mai University, Chiang Mai

This study was undertaken to evaluate the incidence and severity of residual lesions in women featuring high-grade squamous intraepithelial lesion (HSIL) histology with endocervical cone margin involvement after the loop electrosurgical excision procedure (LEEP). The medical records of women undergoing LEEP at Chiang Mai University Hospital between October 2004 and February 2006 were retrospectively reviewed and 74 cases were identified. Nineteen women were excluded because of loss to follow-up. The remaining 4 were referred to other hospitals and 2 declined re-excision, leaving a study population of 55 women for analysis. Mean age \pm SD of the patients was 48.5 \pm 8.9 years. Residual lesions were noted in 26 (47.3%, 95% CI=33.7 to 61.2). Four (7.3%) had unrecognized invasive cervical carcinoma in subsequent specimens. In conclusion, approximately half of women with positive endocervical cone margins after LEEP for HSIL histology have residual disease. Repeat diagnostic excision is recommended for evaluation of lesions and severity.

Published in: *Asian Pac J Cancer Prev* 2006; 7(3): 463-6

WOMEN IN A REGION WITH HIGH INCIDENCE OF CERVICAL CANCER WARRANT IMMEDIATE COLPOSCOPY FOR LOW-GRADE SQUAMOUS INTRAEPITHELIAL LESION ON CERVICAL CYTOLOGY.

Chailert Phongnarisorn¹, Jatupol Srisomboon¹, Sumalee Siriaungkul², Surapan Khunamornpong², Prapaporn Suprasert¹, Kittipat Charoenkwan¹, Chalong Cheewakriangkrai¹, Sitthicha Siriaree¹, Kamonwan Saphan¹

¹ Department of Obstetrics and Gynecology,

² Department of Pathology, Faculty of Medicine, Chiang Mai University,

The objective of this study was to determine the prevalence of high-grade histologic diagnoses in women who had low-grade squamous intraepithelial lesion (LSIL) on Pap smear in an area with high incidence of cervical cancer. We conducted a retrospective study of 220 women with LSIL cytology undergoing colposcopic examination in Chiang Mai University Hospital between January 1999 and July 2004. The histologic diagnoses, obtained from colposcopically directed biopsy or electrical loop excision after initial colposcopy, showed that 80 (36.4%) women had histologically confirmed high-grade lesions and 11 (5%) women had microinvasive (9) and frankly invasive (2) carcinomas. Overall, 41.5% of women with LSIL on Pap smear had significant underlying lesions, requiring appropriate treatment. In conclusion, in the region with high incidence of cervical cancer, women with LSIL cytology are at increased risk of having underlying high-grade lesions and invasive cancer. Immediate referral for colposcopy is warranted.

Published in: *Int J Gynecol Cancer* 2006; 16(4): 1565-8.

HISTOPATHOLOGICAL OUTCOMES OF WOMEN WITH SQUAMOUS CELL CARCINOMA ON CERVICAL CYTOLOGY.

Kittipat Charoenkwan¹, Jatupol Srisomboon¹, Prapaporn Suprasert¹, Sumalee Siriaungkul², Surapan Khunamornpong².

¹ Department of Obstetrics and Gynecology,

² Department of Pathology, Faculty of Medicine, Chiang Mai University

The aim of this study was to determine the final histopathological outcome for women with a squamous cell carcinoma (SCCA) diagnosed by cervical cytology. The medical records and computerized colposcopic database of patients with SCCA on cytology who underwent colposcopy at Chiang Mai University Hospital between February 2003 and December 2005 were reviewed and 48 women with SCCA cytology were identified (mean age 50 years, range 31-73). Nineteen (39.6%) women were postmenopausal. Unsatisfactory colposcopy was noted in 42 (87.5%). Thirty one (64.6%) of the patients had a final pathological diagnosis of high-grade squamous intraepithelial lesions (HGSIL), whereas only 16 (33.3%) had invasive cancer. The remaining one patient had a low-grade squamous intraepithelial lesion. Sensitivity and specificity of colposcopic examination for predicting invasive cancer was 50% and 78%, respectively. In conclusion, most women with a SCCA on cervical cytology have high-grade cervical lesions on final pathology, with only one third demonstrating invasive cancer. The loop electrosurgical excision procedure (LEEP) remains an important measure for combined treatment and diagnosis.

Published in: *Asian Pac J Cancer Prev* 2006; 7(3): 403-6

NERVE-SPARING CLASS III RADICAL HYSTERECTOMY: A MODIFIED TECHNIQUE TO SPARE THE PELVIC AUTONOMIC NERVES WITHOUT COMPROMISING RADICALITY.

Kittipat Charoenkwan, Jatupol Srisomboon, Prapaporn Suprasert, Charuwan Tantipalakov, Chumnan Kietpeerakool

Department of Obstetrics and Gynecology, Faculty of Medicine, Chiang Mai University

The objectives were to describe our nerve-sparing class III radical hysterectomy technique and assess the feasibility and safety of the procedure as well as its impact on voiding function. From January to August 2005, 21 consecutive patients with FIGO stage IB-IIA cervical cancer and 1 patient with clinical stage II endometrial cancer underwent nerve-sparing radical hysterectomy with systematic pelvic lymphadenectomy. The transurethral catheter was removed on the seventh postoperative day. Then intermittent self-catheterization was performed and post-void residual urine volume (PVR) was recorded. The nerve-sparing procedure was completed successfully and safely in all of the patients. Eight (36%) and 6 (27%) patients had the PVR of < 100 ml and < 50 ml respectively at the initial removal of the catheter. On the fourteenth day, 82% and 77% of the patients had the PVR of < 100 ml and < 50 ml, respectively. The mean duration before the PVR became < 50 ml was 11.27 (5-26) days. In conclusion, the technique described in this preliminary study appears safe, adequate, and feasible in our population with satisfactory recovery of voiding function. A larger comparative study is needed on long-term urinary, bowel, and sexual function as well as recurrence and survival.

Published in: Int J Gynecol Cancer 2006; 16(4): 1705-12

BENEFIT OF ELECTROCARDIOGRAPHY DURING FRONT-LINE COMBINATION PACLITAXEL AND CARBOPLATIN CHEMOTHERAPY FOR EPITHELIAL OVARIAN CANCER.

Chumnan Kietpeerakool¹, Jitima Tiyayon¹, Prapaporn Suprasert¹, Runsrut Kanjanavanit², Jatupol Srisomboon¹

¹ Department of Obstetrics and Gynecology,

² Department of Medicine, Faculty of Medicine, Chiang Mai University

OBJECTIVE : To evaluate the patterns of electrocardiography (ECG), cardiac risk factors and its clinical consequence in women with epithelial ovarian cancer (EOC) who received paclitaxel and carboplatin (PC) as front line chemotherapy.

MATERIAL AND METHOD: The medical records and electrocardiographic data of women with EOC who received paclitaxel (175 mg/min²) and carboplatin (AUC=5) every 3 weeks at Chiang Mai University Hospital between January 2000 and December 2004 were reviewed for cardiac risk factors and clinical consequence. **RESULTS:** Among 79 women receiving PC for EOC, 43 (54.4%) had cardiac risk factors. Seventy (88.6%) women had normal ECG, the remaining nine had sinus tachycardia (5), bundle branch block (2), mild T inversion (1), and Wolff-Parkinson-White syndrome (1) before the first course of chemotherapy. Among 70 women with normal initial ECG 8 (11.4%) had sinus tachycardia, one (1.4%) had early depolarization, two (2.9%) had sinus bradycardia and three (4.3%) had sinus arrhythmia in subsequent ECG. All these cardiac disturbances were asymptomatic and needed no intervention, indicating grade I toxicity. The odds ratio of developing abnormal ECG in women with cardiac risk factor was 1.24 (95% CI = 0.33 to 4.64, p = 0.77). Among nine patients with abnormal ECG before the first course of PC, six (66.7%) had subsequent abnormal ECG but all were asymptomatic and no worsening of abnormal ECG pattern was noted.

CONCLUSION: Although paclitaxel and carboplatin chemotherapy could induce abnormal ECG in women with either normal or abnormal prior ECG, its consequence was of no clinical significance. Therefore, the benefit of ECG before each treatment course was theoretically limited.

Published in: J Med Assoc Thai 2006;89(11):1805-10.

PATHOLOGY SLIDE REVIEW IS MANDATORY BEFORE PLANNING TREATMENT FOR REFERRAL PATIENTS WITH GYNECOLOGIC CANCER

Chumnan Kietpeerakool¹, Bancha Changkasiri¹, Surapan Khunamornpong², Sumalee Sisiaungkul², Jatupol Srisomboon¹

¹ Department of Obstetrics and Gynecology,

² Department of Pathology, Faculty of medicine, Chiang Mai University, Chiang Mai

Objective: To evaluate the impact of a pathology slide review on the management of gynecologic cancer patients.

Materials and methods: The medical records of referral patients, with the original pathologic diagnosis of gynecologic cancer treated at Chiang Mai University Hospital between January 2002 and May 2003, were retrospectively reviewed.

Results: During the study period, 402 cases were available for analysis. The slides reviewed were obtained from the following organs: cervix (305), ovary (33), endometrium (32), vulva (14), and others (17). In comparison between the diagnosis after slide review and the original referral diagnosis, the minor discrepancy rate was 41.5% (95% CI: 36.7-46.5%). The major discrepancy rate was 9.0% (95% CI: 6.3-12.2%) and the most common clinical consequence was the modification of planned surgery. There was no major diagnostic discrepancy in the vulva specimens. Among patients with cervical neoplasia, there was a significantly lower rate of major discrepancy among patients with gross lesion than among those without (P < 0.001).

Conclusion: Major diagnostic discrepancies were identified in 9% of referral patients with gynecologic cancer. A pathology slide review is strongly recommended before planning treatment to improve the quality of patient care.

Published in: Asia-Pacific J Clin Oncol 2006; 2: 104-8

CAN ADENOCARCINOMA IN SITU OF THE UTERINE CERVIX BE PREDICTED BEFORE CERVICAL CONIZATION?

Chumnan Kietpeerakool¹, Jatupol Srisomboon¹, Wiboonwan Prompittayarat¹, Panida Kanjanavaha², Ream Peuwsai², Chudsuda Dheerakul².

¹ Department of Obstetrics and Gynecology,

² Obstetrics and Gynecologic Emergency Room Unit, Faculty of Medicine, Chiang Mai University, Chiang Mai

This study was undertaken to determine the effectiveness of Papanicolaou (Pap) smear, colposcopically-directed biopsy (CDB), and endocervical curettage (ECC) in preconization detecting adenocarcinoma in situ (AIS) of the uterine cervix. Women, whose cervical conization specimens contained adenocarcinoma in situ without any associated invasive lesion at Chiang Mai University Hospital between March 1998 and March 2006, were reviewed. During the study period, fifty-one women who matched the study inclusion were identified. Glandular abnormality was detected by Pap smear in 22 women (43.1%). Among 29 women with squamous lesion on Pap smear, 9 additional glandular abnormalities were detected on CDB and/or ECC. In total, a glandular abnormality was suspected in 31 women (60.8%) preoperatively. According to the histological type of AIS, a glandular abnormality suspected from preoperative evaluation was noted in 20 women (70.4%) who had pure AIS. Among women with mixed AIS/HSIL, only 12 women (50.0%) had preoperative evaluation, suggesting a glandular abnormality. These data demonstrate that the sensitivity of combining Pap smear, CDB and/or ECC in detecting glandular lesions before conization containing AIS appears to be suboptimal. Further study concerning the improvement of detecting AIS before conization is warranted to select the most appropriate diagnostic conization method for such lesions.

Published in: Asian Pac J Cancer Prev 2006; 7(4):

SAFETY OF THE LOOP ELECTROSURGICAL EXCISION PROCEDURE IN WOMEN WITH EARLY INVASIVE CERVICAL CARCINOMA.

Chumnan Kietpeerakool, Jatupol Srisomboon.

Department of Obstetrics and Gynecology, Faculty of Medicine, Chiang Mai University, Chiang Mai 50200, Thailand

No abstract

Published in: Int J Gynaecol Obstet 2006; 95(1): 54-5.**OUTCOMES OF LOOP ELECTROSURGICAL EXCISION PROCEDURE FOR CERVICAL NEOPLASIA IN HIV-INFECTED WOMEN**

Chumnan Kietpeerakool, Jatupol Srisomboon, Prapaporn Suprasert, Chailert Phongnarisorn, Kittipat Charoenkwan, Sitthicha Siriaree, Chalong Cheewakriangkrai, Charuwan Tantipalakov, Aree Pantusart.

Department of Obstetrics and Gynecology, faculty of Medicine, Chiang Mai University, Chiang Mai

The objective of this study was to evaluate the treatment outcomes and complications in HIV-infected women undergoing loop electrosurgical excision procedure (LEEP) for cervical neoplasia. The medical record of 60 evaluable HIV-infected women who had abnormal Pap smear and underwent LEEP following colposcopy at Chiang Mai University Hospital between May 1998 and June 2004 were reviewed. Thirty-one (51.7%) had associated genital infection at screening. Twenty-five (41.7%) had opportunistic infection but only 18(30.0%) were treated with antiretroviral therapy. The most common abnormal Pap smear was HSIL (46.7%), followed by LSIL (40.0%). Forty (66.7%) women had clear surgical margins after LEEP. Only 1(1.7%) woman had severe intraoperative hemorrhage. Early and late postoperative hemorrhage were noted in 3 (5%) women of each period. Localized infection of the cervix was detected in 7 (11.7%) women. Two (3.3%) women developed cervical stenosis at 6 month after LEEP. There was no significant difference in overall complications between HIV-infected women and the control group (P=0.24). Among 60 HIV-infected women, no statistical difference in the rate of margins involvement (P=1.00) and complications (P=0.85) could be demonstrated between HIV-infected women who received antiretroviral therapy and those who did not. Disease free rate at 6 and 12 months were 97.1% and 88%, respectively. These data demonstrated that LEEP appears to be safe and effective in HIV-infected women.

Key words: outcomes, loop electrosurgical excision procedure, HIV-infected women, complications, cervical neoplasia**Published in:** Int J Gynecol Cancer 2006; 16(3): 1082-8.**COMPLICATIONS OF LOOP ELECTROSURGICAL EXCISION PROCEDURE FOR CERVICAL NEOPLASIA: A PROSPECTIVE STUDY**

Chumnan Kietpeerakool, Jatupol Srisomboon, Apichart Khobjai, Aunchalee Chandacham, Umpawan Tucksinsook

Department of Obstetrics and Gynecology, Faculty of Medicine, Chiang Mai University, Chiang Mai

Objectives: To evaluate the complications following loop electrosurgical excision procedure (LEEP) for diagnosis and treatment of cervical neoplasia.**Material and Method:** Patients with abnormal cervical cytology who underwent LEEP at Chiang Mai University Hospital between November 2004 and June 2005 were prospectively evaluated for complications.**Results:** During the study periods, 206 patients underwent cervical loop excision for a total of 226 procedures. The mean age of the patients was 41 years (range, 26-72 years). Sixty (29.1%) women were menopausal. The most common abnormal cervical cytology was HSIL (56.3%) followed by LSIL (12.1%). Twenty-five (11.2%) patients received re-excision for positive margin after the first procedure. Intraoperative hemorrhage occurred in 7.9% of the procedures. Early and late postoperative hemorrhage occurred in 0.4% and 2.6% of the procedures, respectively. Eight (3.5%) had postoperative infections and were cured with oral antibiotics. By logistic regression analysis, there was no significant correlation between age, menopausal status, HIV status, re-excision procedure, final histopathology, cone dimension and the complications of LEEP.**Conclusion:** Loop electrosurgical excision procedure is safe for evaluation and treatment of cervical neoplasia with an acceptable and manageable surgical morbidity.**Keywords :** Loop electrosurgical excision procedure, complication, cervical neoplasia.**Published in :** J Med Assoc Thai. 2006 ;89(5):583-7.**CLINICOPATHOLOGIC ANALYSIS OF WOMEN WITH SYNCHRONOUS PRIMARY CARCINOMAS OF THE ENDOMETRIUM AND OVARY: 10 - YEAR EXPERIENCE FROM CHIANG MAI UNIVERSITY HOSPITAL.**Jiraprapa Natee¹, Chumnan Kietpeerakool¹, Jatupol Srisomboon¹, Surapan Khunamornpong², Prapaporn Suprasert¹, Chailert Phongnarisorn¹, Chalong Cheewakriangkrai¹, Sitthicha Siriaree¹, Kittipat Charoenkwan¹, Aree Pantusart¹.¹ Department of Obstetrics and Gynecology,² Department of Pathology, Faculty of Medicine, Chiang Mai University, Chiang Mai.

The aim of this study was to analyze the clinicopathologic features and survival outcomes of women with synchronous primary carcinomas of the endometrium and ovary that were treated at Chiang Mai University Hospital between January 1995 and December 2004. During the study period, 43 women with such tumors were identified. These carcinomas accounted for 0.58% (95%CI=0.42-0.79%) of all gynecologic malignancies. Median age at diagnosis was 49 years (range: 34-60 years). Median body mass index (BMI) was 21.6 kg/m² (range: 15.5-27.7). The majority of women (65%) were premenopausal. The most common presenting symptom was abnormal uterine bleeding (42%), followed by a pelvic mass (30%). Twenty-seven (62.8%, 95%CI= 46.7-77.0%) women had concordant endometrioid carcinomas of the endometrium and ovary. Five (11.6%) women experienced tumor recurrence with median follow up 39 months (range: 1-85 months). The overall 5-year survival was 85.2%. There was no significant difference in survival outcomes among the women who had endometrioid/endometrioid histology and those who had other histological subtypes (P=0.674). In conclusion, synchronous primary carcinomas of the endometrium and ovary, although uncommon, should be considered in differential diagnosis in premenopausal women presenting with abnormal uterine bleeding and ovarian tumors. The prognosis of patients with these tumors appears excellent.

Published in: Asian Pac J Cancer Prev 2006; 7(2): 234-8.

NORTHERN THAI WOMEN WITH HIGH GRADE SQUAMOUS INTRAEPITHELIAL LESION ON CERVICAL CYTOLOGY HAVE HIGH PREVALENCE OF UNDERLYING INVASIVE CARCINOMA.

Nathavut Kantathavorn¹, Chailert Phongnarisorn¹, Jatupol Srisomboon¹, Prapaporn Suprasert¹, Sumalee Siriaunkgul², Surapan Khunamornpong², Kanchana Nimmanahaeminda².

¹ Department of Obstetrics and Gynecology, ² Department of Pathology, Faculty of Medicine, Chiang Mai University, Chiang Mai

The aim of this study was to determine the underlying pathology of women with high grade squamous intraepithelial lesion (HSIL) on cervical cytology. A total of 681 women with HSIL cytology undergoing colposcopic examination at Chiang Mai University Hospital (CMUH) between January 2000 and December 2005 were evaluated for the underlying cervical pathology. The final pathology was diagnosed from the most severe lesions obtained by punch biopsy, loop electrosurgical procedure, cold knife conization or hysterectomy. Underlying high grade cervical lesions including cervical intraepithelial neoplasia grade 2, 3 and adenocarcinoma in situ were noted in 502 (73.7%) women. Invasive cervical carcinoma was identified in 141 (20.7%). The remaining 38 (5.6%) had either low grade or no intraepithelial lesions. No significant difference in the prevalence of underlying high grade and invasive lesions was noted between women with cytologic diagnosis of HSIL from CMUH and other hospitals. In conclusion, northern Thai women with HSIL cytology are at significant risk of having underlying severe cervical lesions, and especially invasive carcinoma which is detected in approximately one-fifth of the cases.

Published in: Asian Pac J Cancer Prev 2006; 7(3): 477-9.

THE CYTOMORPHOLOGIC COMPARISON BETWEEN REHYDRATED AIRDRIED AND CONVENTIONAL WET-FIXED PAP SMEARS.

Krit Jaiwong¹, Kanchana Nimmanahaeminda², Sitthicha Siriaree¹, Surapan Khunamornpong².

¹ Department of Obstetrics and Gynecology,

² Department of Pathology, Faculty of Medicine, Chiang Mai University, Chiang Mai

OBJECTIVE: To compare the cytomorphologic quality of the cervical (Pap) smears between two fixation techniques, rehydration of air-dried smears (AD) versus wet fixation (WF).

MATERIAL AND METHOD: Paired-cervical smears (AD and WF) from 172 women who underwent cervical cytology screening at Chiang Mai University Hospital between August 2004 and September 2004 were prospectively evaluated for the cytologic parameters and the staining qualities. **RESULTS:** The mean age of the 172 women was 41.7 years +/-2 SD 18.1), 27 women (15.7%) were postmenopausal. Absence of red blood cells in the smear background was significantly more frequent in AD smears than in WF specimens ($p = 0.0006$). Air-drying artifact was more frequent in AD smears compared to those of WF ($p = 0.036$) but was of only mild degree in all cases. There was no significant difference between AD and WF smears in the cytoplasmic quality including distinctness of cell border ($p = 0.30$) and satisfactory staining ($p = 0.054$). For the nuclear morphology, there was no significant difference between both fixation techniques in the distinctness of nuclear border ($p = 0.26$) and chromatin crispness ($p = 0.23$) of the endocervical nuclei. In squamous nuclei, AD smears had higher frequency of indistinct nuclear border and hazy chromatin compared to WF smears ($p = 0.003$ each). However, these were observed in only mild degree and did not affect the cytologic interpretation.

CONCLUSION: The quality of AD smears was slightly inferior to WF smears but was still satisfactory for cervical cytology. AD technique may be acceptable as an alternative to wet fixation in cytologic cervical cancer screening.

Published in: J Med Assoc Thai 2006; 89(11): 1811-6.

PRIMARY AND METASTATIC MUCINOUS ADENOCARCINOMAS OF THE OVARY: EVALUATION OF THE DIAGNOSTIC APPROACH USING TUMOR SIZE AND LATERALITY.

Surapan Khunamornpong¹, Prapaporn Suprasert², Jongkolnee Settakorn¹, Suwalee Pojchamarnwiputh¹, Wittanee NaChiangmai¹, Sumalee Siriaunkgul¹

¹ Department of Pathology, ² Department of Obstetrics and Gynecology, Faculty of Medicine, Chiang Mai University

OBJECTIVE: To evaluate the usefulness of the recently proposed algorithm (Seidman JD, Kurman RJ, Ronnett BM. Primary and metastatic mucinous adenocarcinomas in the ovaries: incidence in routine practice with a new approach to improve intraoperative diagnosis. *Am J Surg Pathol* 2003; 27: 985-93 [5]) that classifies mucinous adenocarcinomas of the ovary as primary when they were unilateral > or =10 cm and as metastatic when they were unilateral <10 cm or bilateral.

METHODS: Malignant ovarian neoplasms, which were resected in Chiang Mai University Hospital between 1992 and 2003, were histologically reviewed. Mucinous adenocarcinomas involving the ovary were identified. The medical records and radiologic materials were reviewed in correlation with the pathologic features to identify the primary site.

RESULTS: There were 74 cases of mucinous adenocarcinomas; 16 were primary ovarian; 52, metastatic; and 6 of indeterminate primary site (primary versus metastatic). Primary mucinous adenocarcinomas had a mean size of 16.4 cm and bilateral involvement in 13%. Metastatic mucinous adenocarcinomas had a mean size of 11.7 cm and bilateral involvement in 77%. Excluding the 6 tumors of indeterminate primary site, the proposed algorithm correctly classified primary and metastatic tumors in 84% of 68 cases. Of 21 unilateral mucinous adenocarcinomas > or =10 cm, 62% were primary ovarian. Of 5 unilateral tumors <10 cm, 80% were metastatic. Of 42 bilateral mucinous adenocarcinomas, 95% were metastatic.

CONCLUSION: The algorithm provided high accuracy in the overall prediction of primary and metastatic mucinous adenocarcinomas of the ovary, with major strength in the identification of metastatic tumors by bilaterality or size <10 cm. However, the prediction of primary mucinous adenocarcinomas by unilaterality and size > or =10 cm was less reliable than previously reported. Due to the overlapping features between primary and metastatic tumors and the higher frequency of the latter, the possibility of metastases should always be borne in mind in the evaluation of mucinous adenocarcinomas of the ovary.

Published in: Gynecol Oncol 2006; 101(1): 152-7.

METASTATIC TUMORS TO THE OVARIES: A STUDY OF 170 CASES IN NORTHERN THAILAND.

Surapan Khunamornpong¹, Prapaporn Suprasert², Wittanee Na Chiangmai¹, Sumalee Siriaunkgul¹

¹ Department of Pathology, ² Department of Obstetrics and Gynecology, Faculty of Medicine, Chiang Mai University, Chiang Mai

The cases of malignant ovarian tumors treated at Chiang Mai University hospital between 1992 and 2003 were histologically reviewed. The medical records, the radiologic findings, and the follow-up outcome in the cases suspicious or diagnostic of metastases were reviewed to confirm the diagnosis and to determine the primary sites. Metastatic tumors accounted for 30% of malignant ovarian tumors. A total of 170 cases of metastatic tumors included 117 cases with nongynecologic origin and 53 cases with gynecologic origin. Nongynecologic metastatic tumors were from large intestine (31%), stomach (14%), intrahepatic bile duct (10%), breast (9%), extrahepatic bile duct/gallbladder (7%), appendix (5%), hematologic tumors (3%), others (4%), and unknown primary site (16%). Metastatic gynecologic tumors were from cervix (53%), corpus (34%), fallopian tube (11%), and gestational trophoblastic disease (2%). The proportion of metastatic tumors to malignant ovarian tumors in northern Thailand was comparable to those of the Western or Japanese studies. However, the distribution of the primary sites was different and was correlated with the cancer incidence in Thai women. The majority of mucin-producing adenocarcinomas involving the ovaries were metastatic tumors.

Published in: Int J Gynecol Cancer 2006; 16 Suppl 1: 132-8.

LIMITED VALUE OF VAGINAL CYTOLOGY IN DETECTING RECURRENT DISEASE AFTER RADICAL HYSTERECTOMY FOR EARLY STAGE CERVICAL CARCINOMA.

Napat Injampa¹, Prapaporn Suprasert¹, Jatupol Srisomboon¹,
Chailert Phongnarisorn¹, Kanchana Nimmanahaeminda²,
Sitthicha Siriaree¹, Kittipat Charoenkwan¹, Chalong Cheewakriangkrai¹,
Chumnan Kietpeerakool¹.

¹Department of Obstetrics and Gynecology, ²Department of Pathology,
Faculty of Medicine, Chiang Mai University,
Chiang Mai 50200, Thailand

The objective of this study was to evaluate the value of vaginal cytology in routine surveillance for recurrent cervical cancer after surgery. We reviewed the medical records of 565 patients with stage IB-IIA cervical cancer who were treated with radical hysterectomy and pelvic lymphadenectomy (RHPL) and attended follow-up at Chiang Mai University Hospital between January 2000 and May 2006. With the median follow-up of 35 months (range 1-76 months), 23 (4%) patients developed recurrence. The interval from surgery to detection of recurrence ranged from 5-61 months with a median of 23 months. Of the total 4,376 vaginal smears, 5 (0.1%) showed abnormal cytology but only 1 had malignant cells and tumor recurrence. The sensitivity and specificity of vaginal cytology for detection of recurrence were 4.3% and 99.3%, respectively. In conclusion, vaginal cytology has limited value in detection of recurrence after RHPL for early-stage cervical cancer.

Published in: Asian Pac J Cancer Prev 2006; 7(4):

**PUBLICATIONS
&
PRESENTATIONS**

2007

Early versus delayed (traditional) oral fluids and food for reducing complications after major abdominal gynaecologic surgery.

Kittipat Charoenkwan¹, Phillipson G², Theraporn Vutyavanich¹.

¹Department of Obstetrics and Gynecology, Faculty of Medicine, Chiang Mai University

²New Zealand Center of Reproductive Medicine, Auckland, New Zealand.

BACKGROUND: Traditionally postoperative oral intake is withheld until the return of bowel function. There has been concern that early oral intake would result in vomiting and severe paralytic ileus with subsequent aspiration pneumonia, wound dehiscence, and anastomotic leakage. However, supporting scientific evidence for this traditional practice is lacking and there are potential benefits from early postoperative oral intake.

OBJECTIVES: To assess the effects of early versus delayed (traditional) initiation of oral intake of food and fluids after major abdominal gynaecologic surgery. **SEARCH STRATEGY:** We searched the Menstrual Disorders & Subfertility Group's Specialised Register of controlled trials, the electronic databases (MEDLINE, EMBASE, CINAHL), the Cochrane Controlled Trials Register, and the citation lists of relevant publications in April 2007.

SELECTION CRITERIA: Randomised controlled trials that compared the effect of early versus delayed initiation of oral intake of food and fluids after major abdominal gynaecologic surgery were considered. Early feeding was defined as having oral intake of fluids or food within the first 24 hours after surgery regardless of the presence or absence of the signs that indicate the return of bowel function and delayed feeding was defined after first 24 hours following surgery and only after clinical signs of resolution of postoperative ileus. **DATA COLLECTION AND ANALYSIS:** Studies considered were assessed for methodological quality criteria for inclusion. For dichotomous data, relative risks and 95% confidence intervals were calculated. Continuous data were examined using weighted mean difference and 95% confidence interval. Heterogeneity between the results of different studies were examined by using the forest plot of a meta-analysis, the statistical tests of homogeneity of 2 x 2 tables and the I(2) value.

MAIN RESULTS: Early commencement of oral fluids and food was associated with: increased nausea (one study, 195 patients; relative risk 1.79, 95% confidence interval 1.19 to 2.71), shorter time to the presence of bowel sound (one study, 195 patients; weighted mean difference -0.5 day, 95% confidence interval -0.84 to -0.16), shorter time to first solid diet (two studies, 301 patients; weighted mean difference -1.47 day, 95% confidence interval -2.26 to -0.68), and a trend toward shorter hospital stay (two studies, 301 patients; weighted mean difference -0.73 day, 95% confidence interval -1.52 to 0.07). The shorter hospital stay with early feeding was also evident in the study that reported length of hospital stay in median (-2 days, 4.0 days in early feeding group and 6.0 days in traditional feeding group). There was no significant difference in postoperative ileus, vomiting, and abdominal distension, time to presence of flatus, time to the first passage of stool, postoperative nasogastric tube placement, febrile morbidity, wound complications, and pneumonia.

AUTHORS' CONCLUSIONS: Early feeding after major abdominal gynaecologic surgery is safe however associated with the increased risk of nausea and a reduced length of hospital stay. Whether to adopt the early feeding approach should be individualised. Further studies should focus on the cost-effectiveness, patient's satisfaction, and other physiological changes.

Published in: Cochrane Database Systematic Review. 2007 Oct 17;(4):CD004508

Efficacy of cisplatin in early stage cervical cancer with a long waiting period for surgery.

Prapaporn Suprasert, Theera Thongsong, Jatupol Srisomboon, Charnchai Chailert.

Department of Obstetrics and Gynecology, Faculty of Medicine, Chiang Mai University

This study was undertaken as a prospective trial to evaluate the efficacy and safety of pre-operative cisplatin for controlling the tumor volume of stage IB-IIA cervical cancer patients whose schedule for radical surgery was longer than 3 weeks. Between June 2004 and July 2005, 42 patients were recruited to enter the study. Seventy-five mg/m² of cisplatin was administered for 1-2 courses. Cervical tumor volume was measured 1 day before chemotherapy and 1 day before the operation by using 3-dimensional ultrasound. Reduction of cervical tumor volume was noted in 76.2% of cases. The clinical stage, gross appearance of the tumor, histology and number of chemotherapy courses did not significantly affect chemo-responsiveness. The incidence of lymph node metastases was 16.3%. One patient experienced severe vomiting which could be controlled by ondansetron antiemetic. No severe hematologic or other non-hematologic toxicities were identified. In conclusion cisplatin is effective and safe for administration in a pre-operative setting for early stage cervical cancer patients whose surgical schedule is delayed more than 3 weeks.

Published in: Asian Pacific Journal of Cancer Prevention. 2007 Jan-Mar;8(1):51-4

Factors affecting residual lesion in women with cervical adenocarcinoma in situ after cone excisional biopsy.

Jatupol Srisomboon¹, Chumnan Kietpeerakool¹, Prapaporn Suprasert¹, Sumalee Siriaunkgul², Surapan Khunamornpong², Wiboonwan Prompittayarat¹.

¹ Department of Obstetrics and Gynecology, ² Department of Pathology, Faculty of Medicine, Chiang Mai University

The objective of this study was undertaken to evaluate the factors affecting residual lesion in women with adenocarcinoma in situ (AIS) on cervical conization specimens. The medical records of women with AIS who had no associated invasive carcinoma after cervical conization and underwent subsequent hysterectomy at Chiang Mai University Hospital were reviewed. During March 1998 and March 2006, 45 women were included for analysis. The mean age was 45.2 years (range, 30-66 years). Thirteen (28.9%) women presented with AIS on Pap smear. Thirty (66.7%) underwent loop electrosurgical excision procedure and the remaining 15 (33.3%) underwent cold-knife conization. Twenty (44.4%) women had mixed lesions of AIS and squamous intraepithelial lesion on cervical specimens. Surgical cone margins were clear in 25 (55.6%) women. Eighteen (40%) and two (4.4%) women had involved and non-evaluable cone margins, respectively. Residual lesion was noted in 14 (31.1%) hysterectomy specimens. There was no residual lesion in women with clear cone margins while 72% and 50% of women with involved and non-evaluable cone margins, had residual lesion, respectively. These differences were statistically significant ($P < 0.001$). No significant association between the ECC results and the residual lesion was noted ($P = 0.29$). In conclusion, approximately one-third of women with AIS on cervical conization have residual lesion on subsequent hysterectomy specimens. Only cone margin status is a significant predictor for residual lesion.

Published in: Asian Pacific Journal of Cancer Prevention. 2007 Apr-Jun;8(2):225-8.

How can the overtreatment rate of "see and treat" approach be reduced in women with high-grade squamous intraepithelial lesion on cervical cytology?

Chumnan Kietpeerakool¹, Jatupol Srisomboon¹, Surapan Khunamornpong², Sumalee Siriaunkgul², Wiratchanee Sukkawattananon¹.

¹ Department of Obstetrics and Gynecology,

² Department of Pathology, Faculty of Medicine, Chiang Mai University

Background: The aim of this study was to determine the incidence and predictors of overtreatment in "see and treat" approach using loop electrosurgical excision procedure (LEEP) in women with high-grade squamous intraepithelial lesion (HSIL) on cervical cytology. The overtreatment was considered when LEEP specimens contained no cervical pathology. Between January 2001 and April 2006, 446 women with HSIL on Pap smear underwent colposcopy followed by LEEP at Chiang Mai University Hospital. Mean age of these patients was 45.6 years with a range of 25-78 years. One hundred and twenty-one (27.1%) women were menopausal. Unsatisfactory colposcopy was observed in 357 (80.0%) women. Of 446 women, histologically-confirmed HSIL, invasive cancer, low-grade squamous intraepithelial lesions, and adenocarcinoma in situ were detected in 330 (74.0%), 76 (17.0%), 9 (2.0%), and 5 (1.1%), respectively. The overtreatment rate on LEEP specimens was noted in 26 women or 5.8% (95% confidence interval [CI] = 3.8 to 8.4) of 446 women. By multivariate analysis, postmenopausal status was the only significant independent predictor of overtreatment with an adjusted odds ratio of 2.89 (95% CI = 1.30 to 6.43, P = 0.009). When postmenopausal women were excluded from analysis, the overtreatment rate was reduced to only 4.0%. In conclusion, "see and treat" approach appears to be an appropriate strategy in managing women with HSIL cytology. The overtreatment rate could be reduced when such policy is limited for premenopausal women.

Published in: Asian Pacific Journal of Cancer Prevention. 2007 Apr-Jun;8(2):206-8.

Routine prophylactic application of Monsel's solution after loop electrosurgical excision procedure of the cervix: is it necessary?

Chumnan Kietpeerakool, Jatupol Srisomboon, Prapaporn Suprasert , Chalong Cheewakriangkrai, Kittipat Charoenkwan, Sitthicha Siriaree.

Department of Obstetrics and Gynecology, Faculty of Medicine, Chiang Mai University

AIM: To determine the benefit of an immediate application of Monsel's solution after loop electrosurgical excision procedure (LEEP) of the cervix for preventing postoperative bleeding.

METHODS: This randomized controlled trial was conducted at Chiang Mai University Hospital, Chang Mai, Thailand. Women who were scheduled for LEEP were approached for participation in the study. The primary endpoint was the duration of uncomplicated vaginal bleeding. Secondary endpoints were the incidence of persistent vaginal bleeding, and postoperative complications including severe early bleeding, severe delayed bleeding and infection.

RESULTS: Between October 2004 and May 2006, 285 women with an abnormal Pap-smear, who had undergone LEEP, were randomly allocated to the Monsel's group (n = 140) or control group (n = 145). The baseline outcomes were similar between the two groups. In the Monsel's group, the duration of uncomplicated vaginal bleeding was significantly shorter ($P < 0.001$) and the occurrence of persistent vaginal bleeding was significantly lower ($P = 0.014$) than in the control group. The occurrence of severe complications that required treatment, including bleeding and infection, was not significantly different between the two groups ($P = 0.379$).

CONCLUSION: An application of Monsel's solution after LEEP appears to reduce the duration of postoperative vaginal bleeding, but does not significantly prevent severe complications. Such practice may not be necessary if adequate hemostasis is achieved using electrical cauterization.

Published in: *Journal of Obstetrics and Gynaecology Research*. 2007 Jun;33(3):299-304.

Cervical intraepithelial neoplasia II-III with endocervical cone margin involvement after cervical loop conization: is there any predictor for residual disease?

Chumnan Kietpeerakool¹, Surapan Khunamornpong², Jatupol Srisomboon¹, Sumalee Siriaunkgul², Prapaporn Suprasert¹.

¹Department of Obstetrics and Gynecology,

²Department of Pathology, Faculty of Medicine, Chiang Mai University

AIM: To determine the clinicopathological predictors for residual disease in women who have had cervical intraepithelial neoplasia (CIN) II-III with endocervical cone margin involvement after loop electrosurgical excision procedure (LEEP).

METHODS: All of the women who had CIN II-III on LEEP specimens with endocervical margin involvement, and underwent subsequent surgical treatment including repeat LEEP or hysterectomy at Chiang Mai University Hospital between May 2003 and June 2006 were reviewed.

RESULTS: During the study period, 85 women who matched the study inclusion were identified. The mean age was 48.6 years. Fifty-two women (61.2%) were postmenopausal. The most common Pap smear before LEEP was high-grade squamous intraepithelial lesion (HSIL) (65.9%), followed by squamous cell carcinoma (21.2%). Twenty-five women (29.4%) had concurrent ectocervical and endocervical cone margin involvement. Residual disease was noted in 44 women (51.8%, 95%CI = 40.7-62.7) of whom six had unrecognized invasive squamous cell carcinoma, while the remaining 38 had CIN II-III. Only extensive endocervical cone margin involvement (3-4 quadrants) was noted as the significantly independent predictor for residual disease (aOR = 14.2, 95% CI = 3.6-55.8; P < 0.001).

CONCLUSION: Extensive endocervical cone margin involvement after LEEP for CIN II-III is a strong predictor for residual disease. Therefore, the number of involved quadrants should be evaluated to plan further management.

Published in: Journal of Obstetrics and Gynaecology Research. 2007 Oct; 33(5):660-4

Appropriate Interval for Repeat Excision in Women Undergoing Prior Loop Electrosurgical Excision Procedure for Cervical Neoplasia

Chumnan Kietpeerakool, Jatupol Srisomboon, Jitima Tiyayon, Irene Ruengkachorn,
Chalong Cheewakriangkrai, Prapaporn Suprasert, Aree Pantusart.
Department of Obstetrics and Gynecology, Faculty of Medicine, Chiang Mai University

The objective of the present study was to evaluate the impact of intervals on the complications and pathological examination in women undergoing a repeat loop electrosurgical excision procedure (LEEP) for cervical neoplasia. During October 2004 and January 2007, 78 women who had undergone repeat LEEP at Chiang Mai University Hospital, were prospectively evaluated. The mean age was 47.5 years (range; 27-69 years). The mean duration of uncomplicated vaginal bleeding was 4.4 days (range; 1-20 days). The occurrence of persistent vaginal bleeding was noted in 9 (11.5%) women. Among 78 women, 2 (2.56%) and 7 (8.97%) experienced intraoperative and postoperative hemorrhage, respectively. Six (7.69%) had postoperative infection. These complications were not significantly different from those observed in women undergoing single LEEP in the same period ($P=0.56$). There was no significant difference in the incidence of perioperative complications and the incidence of non-evaluable cone margins among women who undergoing repeat LEEP within 4-6 weeks, between 6-8 weeks, and more than 8 weeks after first LEEP. In conclusion, repeat LEEP could be safely performed 4-12 weeks after the first procedure without any impact on pathological specimen examination.

Published in: Asian Pacific Journal of Cancer Prevention 2007; 8(3) 379-382

Underlying Pathology of Women with Atypical Squamous Cells, Cannot Exclude High-Grade Squamous Intraepithelial Lesion (ASC-H) Smears in a Region with a High Incidence of Cervical Cancer

Chumnan Kietpeerakool¹, Jatupol Srisomboon¹, Charuwan Tantipalakovorn¹, Prapaporn Suprasert¹, Surapan Khunamornpong², Kanchana Nimmanhaeminda², Sumalee Siriaunkgul²

¹ Department of Obstetrics and Gynecology,

² Department of Pathology, Faculty of Medicine, Chiang Mai University

Aim: To evaluate the histopathology of women, who had atypical squamous cells, cannot exclude high-grade squamous intraepithelial lesions (ASC-H) on cervical cytology in a region with high incidence of cervical cancer.

Methods: This study was conducted at Chiang Mai University Hospital, Chiang Mai, Thailand. All women with ASC-H, who had undergone colposcopic and histopathologic evaluation between October 2004 and January 2007, were recruited. Similar cohorts with other squamous cell abnormalities on a Pap smear, who had undergone colposcopy during the same period, were included as comparative groups.

Results: During the study period, 85 women who had ASC-H smears underwent colposcopic and histopathologic evaluation. The mean age was 45.3 years (range, 20-64 years). The histopathologic results of these 85 women were as follows: CIN II-III, 52 (61.2%); invasive cancer, 7(8.2%); CIN I, 6 (7.1%); and no lesions, 20 (23.5%). The incidence of underlying CIN II or higher in an ASC-H smear (69.4%) was intermediate between ASC-US (22.7%), LSIL (44.7%) and HSIL (90.5%) smears. There was no statistically significant difference in the incidence of CIN II or higher between women who were 40 years old or more and those who were younger (68.7% and 71.4%, respectively, P=0.81), or between premenopausal and postmenopausal women (71.4% and 63.6%, respectively, P=0.49).

Conclusion: Reporting ASC-H cytology in our population is strongly associated with significant cervical pathology, particularly invasive cancer that is possibly at a rate higher than previously reported. Women who have ASC-H smears should therefore be referred for immediate colposcopy regardless of age and menopausal status.

Published in: Journal of Obstetrics and Gynaecology Research 2008: (in press)

Lymphovascular space invasion as a prognostic determinant in uterine cancer

Chalong Cheewakriangkrai¹, Kannika Panggid¹, Sumalee Siriaungkul², Surapan Khunamornpong², Prapaporn Suprasert¹, Jatupol Srisomboon¹

¹Department of Obstetrics and Gynecology,

²Department of Pathology, Faculty of Medicine, Chiang Mai University, Chiang Mai

The objective of this study was to evaluate the clinical significance of lymphovascular space invasion (LVSI) in patients with uterine cancer in terms of lymph node metastasis, recurrence and survival rate. A total of 190 patients with newly diagnosed uterine cancer who underwent total abdominal hysterectomy (TAH), bilateral salpingo-oophorectomy (BSO), omentectomy, peritoneal washing or ascitic fluid collection, and pelvic/paraortic lymph node sampling at Chiang Mai University Hospital between January 1999 and December 2004 were evaluated. All medical records and histopathologic slides were retrospectively reviewed to determine the relationship between LVSI and clinicopathological characteristics. LVSI was present in 79 patients (42%) and significantly correlated with lymph node metastasis ($p < 0.001$), BMI $< 25 \text{ kg/m}^2$ ($p < 0.001$), advanced FIGO stage ($p < 0.001$), poor histologic grade ($p < 0.001$), and deep uterine invasion ($p < 0.001$). Patients with LVSI, when stratified by FIGO stage, also had a significant lower 5-year survival rate. For those who had disease recurrences, LVSI and histologic grade were found to be independent prognostic factors in a multivariate analysis. LVSI was one of the prognostic determinants for disease recurrence and associated with poor survival in patients with uterine cancer.

Published in: Asian Pacific Journal of Cancer Prevention 2007; 8(3): 363-36

Sexual function after radical hysterectomy for early-stage cervical cancer.

Jantana Jongpipat, Kittipat Charoenkwan.

Department of Obstetrics and Gynecology, Faculty of Medicine, Chiang Mai University

INTRODUCTION: For early-stage cervical cancer, treatment with radical surgery appears effective with excellent survival. However, the treatment is associated with significant morbidities. Sexual dysfunction is the leading cause of symptom-induced distress after the treatments for early-stage cervical cancer. There has been no study that evaluates the effect of surgical treatments for cervical cancer on sexual function in the Eastern/Asian patients.

AIM: To examine the effect of radical hysterectomy on postoperative sexual function in women with early-stage cervical cancer.

MAIN OUTCOME MEASURES. The visual analog score on seven aspects of sexual function: overall satisfaction with sexual intercourse, sexual desire, vaginal lubrication, vaginal elasticity, orgasmic satisfaction, patient-perceived partner satisfaction, and associated anxiety.

METHODS: Thirty patients with International Federation of Gynecology and Obstetrics (FIGO) stage IB cervical cancer undergoing radical hysterectomy and pelvic lymphadenectomy at the Chiang Mai University Hospital were recruited. All of the patients were interviewed by questionnaire on sexual function at preoperative hospital admission and then at 3 and 6 months after surgery. Relevant clinical and pathologic data were also prospectively collected. **RESULTS:** The mean age of the patients was 45.3 +/- 6 years (range 28-59). Seven (23%) of the patients were postmenopausal. Almost all (97%) had FIGO stage IB1 disease. Fourteen (46.7%) patients had bilateral salpingo-oophorectomy, and one (3.3%) patient had unilateral salpingo-oophorectomy. Squamous-cell carcinoma and adenocarcinoma were diagnosed in 83% and 17%, respectively. At 3 and 6 months after operation, 63% and 93% of the patients had sexual intercourse, respectively. Considering the overall sexual satisfaction, the patient-rated visual analog score appeared comparable at preoperative and 6-month postoperative assessment (4.2 +/- 2.5 vs. 3.6 +/- 2.2, $P > 0.05$). **CONCLUSION:** Radical hysterectomy using the current technique has a minimal short-term impact on sexual function. A comparative study with long-term follow-up would provide clearer information on permanent effect of radical hysterectomy on sexual function.

Published in: Journal of Sexual Medicine. 2007 Nov;4(6):1659-65.

Prevalence and characteristics of late postoperative voiding dysfunction in early-stage cervical cancer patients treated with radical hysterectomy

Kittipat Charoenkwan, Savitree Pranpanas

Department of Obstetrics and Gynecology, Faculty of Medicine, Chiang Mai University, Chiang Mai

Objective: To examine prevalence and characteristics of voiding dysfunction following radical hysterectomy for early-stage cervical cancer

Methods: One hundred-eighty seven patients with FIGO stage IA2-IIA cervical cancer who underwent class II-III radical hysterectomy with systematic pelvic lymphadenectomy between January 1, 2002 and June 31, 2005 were interviewed with questionnaire on voiding function. Medical records were also reviewed for operative and pathologic data.

Results: The prevalence of symptomatic bladder dysfunction was 25.1%. There was no statistically significant association between rates of bladder dysfunction and all examined clinical/operative factors. The most common pattern of bladder dysfunction were incomplete emptying in 25 (13.4%) and urgency and nocturia in 21 (11.2%) each.

Conclusion: Voiding dysfunction is a common and clinically significant long-term complication following radical hysterectomy. The pattern of dysfunction reflects combined surgical disruption of both parasympathetic and sympathetic innervations of the pelvis.

Published in: Asian Pacific Journal of Cancer Prevention 2007; 8(3): 387-389.

Surgical morbidity associated with total laparoscopic hysterectomy in women with prior diagnostic excision of the cervix.

Chailert Phongnarisorn, Jatupol Srisomboon.

Department of Obstetrics & Gynecology, Faculty of Medicine, Chiang Mai University, Chiang Mai

AIMS: To evaluate the feasibility, safety, and complications of total laparoscopic hysterectomy (TLH) in women undergoing prior diagnostic excision of the cervix.

METHODS: A retrospective study (Canadian Task Force classification II-2) was conducted in a tertiary care university hospital. The medical records of women undergoing TLH between June 2003 and September 2004 were reviewed.

RESULTS: Twenty-six women with stage IA1 cervical cancer (19) and persistent high grade cervical neoplasia (7) underwent TLH after diagnostic cervical excision. The operation was successfully performed in all cases without conversion to laparotomy. The mean age of the patients was 47.0 +/- 8.64 years (95% CI 43.5-50.5). 19 patients had previous loop electrosurgical excision procedure (LEEP), one had cold knife conization. Six patients underwent repeated LEEP for positive endocervical margin. The mean operating time was 253.0 +/- 66.7 min (95% CI 226.0-279.9). The median blood loss was 300 mL (range 50-1000 mL). Only one patient needed 1 unit of blood transfusion. The median post-operative hospital stay was 3 days (range 2-6 days). All hysterectomy specimens had negative surgical margins. Two patients had major complications, one with bladder injury requiring laparoscopic repair. The remaining one had ureteral injury detected 9 days after the operation requiring subsequent ureteroneocystostomy. Both complications occurred in the first four cases of this series. No significant morbidity was noted in 2 years of follow-up. **CONCLUSION:** TLH appears to be feasible and safe in patients with prior diagnostic excision of the cervix. Careful separation of the bladder from the cervix and identification of both ureters are recommended to minimize morbidity associated with this operation.

Published in: Journal of Obstetrics and Gynaecology Research. 2007 Aug;33(4):519-23

Effects of gel lubricant on cervical cytology

Kittipat Charoenkwan¹, Karnchana Nimmanahaeminda²,
Surapan Khunamornpong², Jatupol Srisomboon¹, Thorner, PS.³

¹ Department of Obstetrics and Gynecology,

² Department of Pathology, Faculty of Medicine, Chiang Mai University

³ Department of Laboratory Medicine, Hospital for Sick Children and University of Toronto, Toronto, Canada

Objective: To determine whether the use of lubricating gel during vaginal speculum examination affected cytological interpretation in the conventional Papanicolaou (Pap) smear.

Methods: Two consecutive cervical smears were obtained from 1,334 patients undergoing Pap smear screening. The first smear ('non-contaminated') was obtained using the routine collection technique. The second smear ('gel-contaminated') was taken after applying a 1 to 1.5 cm ribbon of lubricating gel onto the external cervical os. Adequacy of Pap smear and discordance in diagnosis between the paired smears were examined.

Results: The proportion of unsatisfactory smears was significantly higher in the gel-contaminated smears, 12.1% versus 1.7% ($p < 0.01$). This difference was consistent across all reproductive groups. For patients who had smears satisfactory for cytological evaluation, the discordance in cytological diagnosis between the gel-contaminated and non-contaminated smears from the same patient was 0.3%.

Conclusion: Lubricating gel contamination of the cervix can adversely affect adequacy and cytological diagnosis in the conventional Pap smear.

Published in: *Acta Cytologica* 2008 (in press).

Poster Presentation

TEN YEARS EXPERIENCE WITH RADICAL HYSTERECTOMY AND PELVIC LYMPHADENECTOMY IN EARLY STAGE CERVICAL CANCER

Prapaporn Suprasert, Chailert Phongnarisorn, Kittipat Charoenkwan , Chalong Cheewakriangkrai, Sitthicha Siriaree, Chumnan Kietpeerakool , Charuwan Tantipalakorn, Jatupol Srisomboon.

Division of Gynecologic Oncology, Department of Obstetrics and Gynecology.
Faculty of Medicine, Chiang Mai University

Background & Aim: Radical hysterectomy and pelvic lymphadenectomy (RHPL) is the main surgical treatment in early stage cervical cancer. With the long experience of this procedure in our institute, we conducted the retrospective study to evaluate the outcome and complication of this treatment.

Methods: The medical record of cervical cancer patients undergoing RHPL at Chiang Mai university hospital between January 1995 and December 2004 were reviewed.

Result: There were 990 available data patients in the study period. The mean age was 43 years old (range 17-79 years). 251 patients had prior diagnostic conization. The maximum tumor size was 6 cm. The most common histology was squamous cell carcinoma (67%) followed by adenocarcinoma (24%). The distribution of FIGO staging were stage IA (9%), stage IB1(60%), stage IB (20%), stage IB2 (4%) and stage IIA (7%). Pelvic node, parametrial and vaginal margin metastases were detected in 16.6%, 8.9% and 3.3% of patients, respectively. 67% of patients underwent RHPL without adjuvant treatment. 62 (6%) patients developed recurrence and 5 died of disease. The estimated -10 year recurrence- free survival was 92%. The most common long term complication was lymphedema which was noted in 7% of patients.

Conclusion: Early stage cervical cancer treated with RHPL has long term favorable outcome with minimal morbidity.

Poster Presented in 15th ESGO, Berlin, German, 31 October, 2007